






## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : <b>A61K 31/415, 31/41, C07D 471/02</b></p>	<p><b>A1</b></p>	<p>(11) International Publication Number: <b>WO 00/15222</b> (43) International Publication Date: 23 March 2000 (23.03.00)</p>
<p>(21) International Application Number: PCT/US99/21070 (22) International Filing Date: 13 September 1999 (13.09.99) (30) Priority Data: 60/100,665 16 September 1998 (16.09.98) US (71) Applicant: BRISTOL-MYERS SQUIBB COMPANY [US/US]; P.O. Box 4000, Princeton, NJ 08543-4000 (US). (72) Inventors: MACOR, John, E.; 258 Kuhl Road East, Flemington, NJ 08822 (US). YU, Guixue; 8 Greene Drive, Lawrenceville, NJ 08648 (US). (74) Agents: DAVIS, Stephen, B. et al.; Bristol-Myers Squibb Company, P.O. Box 4000, Princeton, NJ 08543-4000 (US).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: FUSED PYRIDINE INHIBITORS OF cGMP PHOSPHODIESTERASE</p> <div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;">  <p>(I)</p> </div> <div style="text-align: center;">  <p>(II)</p> </div> <div style="text-align: center;">  <p>(III)</p> </div> </div> <p>(57) Abstract</p> <p>Compounds of the formulas (I), (II) and (III) are useful as inhibitors of cGMP PDE, especially type V.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## FUSED PYRIDINE INHIBITORS OF cGMP PHOSPHODIESTERASE

5

Field of the Invention

The present invention relates to fused pyridine compounds, to methods of using such compounds in the treatment of cGMP-associated conditions such as erectile dysfunction, and to pharmaceutical compositions containing such compounds.

Background of the Invention

15       Erectile dysfunction is the inability to obtain and maintain a penile erection sufficient for sexual intercourse or other sexual expression. A number of factors can place an individual at risk for this disorder, for example, trauma, pelvic surgery, hypercholesterolemia, ischemic heart disease, peripheral vascular disease, chronic renal failure, diabetes, or the use of medicaments such as certain antihypertensive medications or digoxin, or illicit drugs, cigarettes or alcohol. Methods for the treatment of erectile

20       dysfunction include the use of vacuum devices and penile implants, as well as the administration of medicaments such as yohimbine, papaverine and apomorphine. Improved methods for the treatment of this disorder are sought, however, as the aforementioned methods do not provide

25       sufficient efficacy, and/or are accompanied by drawbacks or side effects such as erosion, pain, priapism or gastrointestinal discomfort.

30       As penile erection is dependent upon the presence of adequate levels of cyclic guanosine 3',5'-monophosphate (cGMP), especially in corpora cavernosa tissue,

35


administration of an inhibitor of a cGMP phosphodiesterase (cGMP PDE) (and particularly, a selective inhibitor of cGMP PDE Type V (cGMP PDE V)) provides a means for achieving and maintaining an erection, and therefore for treating erectile dysfunction. See Trigo-Rocha et al., "Nitric Oxide and cGMP: mediators of pelvic nerve-stimulated erection in dogs," *Am. J. Physiol.*, Vol. 264 (Feb. 1993); Bowman et al., "Cyclic GMP mediates neurogenic relaxation in the bovine retractor penis muscle," *Br. J. Pharmac.*, 81, 665-674 (1984); and Rajfer et al., "Nitric Oxide as a Mediator of Relaxation of the Corpus Cavernosum in Response to Nonadrenergic, Noncholinergic Neurotransmission," *New England J. Med.*, 326, 2, 90-94 (Jan. 1992). Sildenafil, for example, has been described as a phosphodiesterase Type V inhibitor useful for the treatment of erectile dysfunction. See *Drugs of the Future*, 22, 138-143 (1997).

The present invention provides novel compounds which are potent and selective inhibitors of cGMP PDE V which may be employed in the treatment of erectile dysfunction. In view of their activity, the present compounds can also be employed in the treatment of other disorders responding to the inhibition of cGMP PDE such as various cardiovascular disorders.


#### Summary of the Invention

This invention is directed to the novel fused pyridine compounds of formulas I and II shown below including pharmaceutically acceptable salts thereof, pharmaceutical compositions containing one or more fused pyridines of formulas I and II, and the use of such compounds as inhibitors of cGMP PDE, especially type V.

(I)



(II)




5

This invention is also directed to the use of the fused pyridine compounds of formula III shown below including pharmaceutically acceptable salts thereof as inhibitors of cGMP PDE, especially type V.

10

(III)




In the above formulas:


15

E<sub>1</sub> is -O-R<sub>1</sub>, -S-R<sub>1</sub>, -NH-A<sub>1</sub>-cycloalkyl, -NH-A<sub>1</sub>-substituted cycloalkyl, -NH-A<sub>1</sub>-heterocyclo, or -NH-A<sub>1</sub>-heteroaryl.





$E_2$  is  $-NH-A_1$ -alkoxy,  $-NH-A_1-CO_2$ alkyl,




- 5  $R_1$  is  $-A_1$ -cycloalkyl,  $-A_1$ -substituted cycloalkyl,  $-A_1$ -alkoxy,  $-A_1-N \begin{matrix} \nearrow R_{15} \\ \searrow R_{16} \end{matrix}$ ,  $-A_1$ -aryl,  $-A_1$ -substituted aryl,  $-A_1$ -heterocyclo, or  $-A_1$ -heteroaryl.



- 10 monocyclic ring  $\begin{matrix} -N \\ \diagup \quad \diagdown \\ (R_{21})_n \quad (R_{22})_m \end{matrix}$ , a fused bicyclic ring



15



5  $A_1$  is an alkylene or substituted alkylene bridge of 1 to 10 carbons.

$Y$  is nitrogen or  $C(R_6)$ .

$Z$  is nitrogen or  $C(R_7)$  with the proviso that at least one of  $Y$  or  $Z$  is nitrogen.

10  $R_3$  is hydrogen, alkyl, cycloalkyl, substituted cycloalkyl, substituted alkyl,  $-A_1$ -aryl,  $-A_1$ -substituted aryl,  $-A_1$ -cycloalkyl, or  $-A_1$ -substituted cycloalkyl.

15  $R_6$  and  $R_7$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl,  $-A_1$ -cycloalkyl,  $-A_1$ -substituted cycloalkyl,  $-A_1$ -aryl,  $A_1$ -substituted aryl,  $-A_1$ -heterocyclo, and  $A_1$ -heteroaryl.

$R_4$  is hydrogen,  $-N(R_{12})(R_{13})$ ,  $-OR_{12}$  or 1- or 3-imidazolyl.

20  $A_2$  is a direct bond, an alkylene or substituted alkylene bridge of 1 to 10 carbons, an alkenyl or substituted alkenyl bridge of 2 to 10 carbons having one or more double bonds, or an alkynyl or substituted alkynyl bridge of 2 to 10 carbons having one or more triple bonds.

25  $R_2$  is cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclo, heteroaryl, cycloalkyl- $A_3$ -cycloalkyl, cycloalkyl- $A_3$ -substituted cycloalkyl, cycloalkyl- $A_3$ -aryl, cycloalkyl- $A_3$ -substituted aryl, cycloalkyl- $A_3$ -heterocyclo, cycloalkyl- $A_3$ -heteroaryl, 30 substituted cycloalkyl- $A_3$ -cycloalkyl, substituted cycloalkyl- $A_3$ -substituted cycloalkyl, substituted cycloalkyl- $A_3$ -aryl, substituted cycloalkyl- $A_3$ -substituted

- aryl, substituted cycloalkyl-A<sub>3</sub>-heterocyclo, substituted cycloalkyl-A<sub>3</sub>-heteroaryl, aryl-A<sub>3</sub>-cycloalkyl, aryl-A<sub>3</sub>-substituted cycloalkyl, aryl-A<sub>3</sub>-aryl, aryl-A<sub>3</sub>-substituted aryl, aryl-A<sub>3</sub>-heterocyclo, aryl-A<sub>3</sub>-heteroaryl, substituted
- 5 aryl-A<sub>3</sub>-cycloalkyl, substituted aryl-A<sub>3</sub>-substituted cycloalkyl, substituted aryl-A<sub>3</sub>-aryl, substituted aryl-A<sub>3</sub>-substituted aryl, substituted aryl-A<sub>3</sub>-heterocyclo, substituted aryl-A<sub>3</sub>-heteroaryl, heterocyclo-A<sub>3</sub>-cycloalkyl, heterocyclo-A<sub>3</sub>-substituted cycloalkyl, heterocyclo-A<sub>3</sub>-
- 10 aryl, heterocyclo-A<sub>3</sub>-substituted aryl, heterocyclo-A<sub>3</sub>-heterocyclo, heterocyclo-A<sub>3</sub>-heteroaryl, heteroaryl-A<sub>3</sub>-cycloalkyl, heteroaryl-A<sub>3</sub>-substituted cycloalkyl, heteroaryl-A<sub>3</sub>-aryl, heteroaryl-A<sub>3</sub>-heterocyclo, heteroaryl-A<sub>3</sub>-heteroaryl, cyano, -OR<sub>9</sub>, -SR<sub>9</sub>, -(C=O)R<sub>9</sub>, -N(R<sub>9</sub>)(R<sub>10</sub>),
- 15 -CO<sub>2</sub>R<sub>9</sub>, -(C=O)N(R<sub>12</sub>)(R<sub>13</sub>), -SO<sub>2</sub>N(R<sub>12</sub>)(R<sub>13</sub>), -NR<sub>11</sub>(C=O)R<sub>19</sub>, -NR<sub>11</sub>(C=O)N(R<sub>12</sub>)(R<sub>13</sub>), -O-(C=O)N(R<sub>12</sub>)(R<sub>13</sub>) provided that A<sub>2</sub> is not a direct bond, -NR<sub>11</sub>CO<sub>2</sub>R<sub>19</sub>, -(C=O)N(R<sub>11</sub>)CH<sub>2</sub>CO<sub>2</sub>R<sub>19</sub>, nitrogen when A<sub>2</sub> is alkynyl ending in a triple bond, or NH when A<sub>2</sub> is alkenyl ending in a double bond.
- 20 R<sub>25</sub> is cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclo, heteroaryl, cycloalkyl-A<sub>3</sub>-cycloalkyl, cycloalkyl-A<sub>3</sub>-substituted cycloalkyl, cycloalkyl-A<sub>3</sub>-aryl, cycloalkyl-A<sub>3</sub>-substituted aryl, cycloalkyl-A<sub>3</sub>-heterocyclo, cycloalkyl-A<sub>3</sub>-heteroaryl,
- 25 substituted cycloalkyl-A<sub>3</sub>-cycloalkyl, substituted cycloalkyl-A<sub>3</sub>-substituted cycloalkyl, substituted cycloalkyl-A<sub>3</sub>-aryl, substituted cycloalkyl-A<sub>3</sub>-substituted aryl, substituted cycloalkyl-A<sub>3</sub>-heterocyclo, substituted cycloalkyl-A<sub>3</sub>-heteroaryl, aryl-A<sub>3</sub>-cycloalkyl, aryl-A<sub>3</sub>-
- 30 substituted cycloalkyl, aryl-A<sub>3</sub>-aryl, aryl-A<sub>3</sub>-substituted aryl, aryl-A<sub>3</sub>-heterocyclo, aryl-A<sub>3</sub>-heteroaryl, substituted aryl-A<sub>3</sub>-cycloalkyl, substituted aryl-A<sub>3</sub>-substituted cycloalkyl, substituted aryl-A<sub>3</sub>-aryl, substituted aryl-A<sub>3</sub>-substituted aryl, substituted aryl-A<sub>3</sub>-
- 35 heterocyclo, substituted aryl-A<sub>3</sub>-heteroaryl, heterocyclo-



- $A_3$ -cycloalkyl, heterocyclo- $A_3$ -substituted cycloalkyl,  
 heterocyclo- $A_3$ -aryl, heterocyclo- $A_3$ -substituted aryl,  
 heterocyclo- $A_3$ -heterocyclo, heterocyclo- $A_3$ -heteroaryl,  
 heteroaryl- $A_3$ -cycloalkyl, heteroaryl- $A_3$ -substituted  
 5 cycloalkyl, heteroaryl- $A_3$ -aryl, heteroaryl- $A_3$ -substituted  
 aryl, heteroaryl- $A_3$ -heterocyclo, heteroaryl- $A_3$ -heteroaryl,  
 cyano,  $-S-R_9$ ,  $-(C=O)R_{11}$ ,  $-CO_2R_{19}$ ,  $-(C=O)N(R_{12})(R_{13})$ ,  
 $-SO_2N(R_{12})R_{13}$ ,  $-NF_9(C=O)R_{10}$ ,  $-NR_{11}(C=O)N(R_{12})(R_{13})$ ,  
 $-O-(C=O)N(R_{12})(R_{13})$  provided that  $A_2$  is not a direct bond,  
 10  $-NR_{11}CO_2R_{19}$ ,  $-(C=O)N(R_{11})CH_2CO_2R_{19}$ , nitrogen when  $A_2$  is  
 alkynyl ending in a triple bond, or NH when  $A_2$  is alkenyl  
 ending in a double bond.


- $A_3$  is a direct bond, an alkylene or substituted  
 alkylene bridge of 1 to 10 carbons, an alkenyl or  
 15 substituted alkenyl bridge of 2 to 10 having one or more  
 double bonds, an alkynyl or substituted alkynyl bridge of  
 2 to 10 carbons having one or more triple bonds,  
 $-(CH_2)_d-O-(CH_2)_e-$ ,  $-(CH_2)_d-S-(CH_2)_e-$ , or  
 $-(CH_2)_d-(C=O)-(CH_2)_e-$ .

- 20  $d$  is zero or an integer from 1 to 6.

$e$  is zero or an integer from 1 to 6.

- $R_5$  is hydrogen, alkyl, substituted alkyl,  
 cycloalkyl, substituted cycloalkyl, aryl,  $-A_1$ -aryl,  
 substituted aryl,  $-A_1$ -substituted aryl, heterocyclo,  $-A_1$ -  
 25 heterocyclo, heteroaryl or  $-A_1$ -heteroaryl.

- $R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$ ,  $R_{13}$ ,  $R_{15}$ ,  $R_{16}$ , and  $R_{19}$  are  
 independently selected from the group consisting of  
 hydrogen, alkyl, substituted alkyl, cycloalkyl,  
 substituted cycloalkyl, aryl, substituted aryl,  
 30 heterocyclo, heteroaryl,  $-A_1$ -cycloalkyl,  $-A_1$ -substituted  
 cycloalkyl,  $-A_1$ -aryl,  $-A_1$ -substituted aryl,  $-A_1$ -heterocyclo  
 and  $-A_1$ -heteroaryl, or  $R_{12}$  and  $R_{13}$  taken together with the  
 N atom to which they are attached represent a heterocyclo  
 ring.



represents a monocyclic heterocyclo or heteroaryl ring of 4 to 8 atoms containing up to 3 additional heteroatoms (up to 2 additional heteroatoms when the ring is 4 atoms) which are selected from one or two oxygen atoms and/or one or two sulfur atoms and/or one, two or three nitrogen atoms.


$R_{21}$  is attached to an available carbon or nitrogen atom and is hydrogen, alkyl, halogen, hydroxy, trifluoromethyl, amino, alkoxy or carboxy.


$R_{22}$  is attached to an available carbon or nitrogen atom and is keto,  $-(C=O)R_{23}$ ,  $-CO_2-R_{23}$ ,  $-NH-(C=O)-R_{23}$ ,  $-N(alkyl)_2$ ,  $-A_1$ -hydroxy,  $-A_1-N(R_9)(R_{10})$ ,  $-A_1$ -alkoxy,  $-A_1$ -carboxy,  $-A_2$ -cycloalkyl,  $-A_2$ -substituted cycloalkyl,  $-A_2$ -aryl,  $-A_2$ -substituted aryl,  $-A_2$ -heterocyclo, or  $-A_2$ -heteroaryl.

$n$  is one or two.

$m$  is zero or one.


$R_{23}$  is alkyl,  $-N(R_9)(R_{10})$ ,  $-A_1$ -hydroxy,  $-A_1-N(R_9)(R_{10})$ ,  $-A_1$ -carboxy,  $-A_2$ -cycloalkyl,  $-A_2$ -substituted cycloalkyl,  $-A_2$ -aryl,  $-A_2$ -substituted aryl,  $-A_2$ -heterocyclo or  $-A_2$ -heteroaryl.




represents a fused bicyclic ring wherein the monocyclic ring  is defined previously and






represents a cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclo or heteroaryl having two carbon atoms in common with the monocyclic ring





represents a spiro ring wherein the

monocyclic ring  is defined previously and  represents a cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclo or heteroaryl ring having a

5 common carbon with the monocyclic ring  .

### Detailed Description of the Invention

10 The following are definitions of terms used in this specification. The initial definition provided for a group or term herein applies to that group or term throughout the present specification, individually or as part of another group, unless otherwise indicated.

15 The term "alkyl" refers to straight or branched chain hydrocarbon groups having 1 to 12 carbon atoms, preferably 1 to 8 carbon atoms. Lower alkyl groups, that is, alkyl groups of 1 to 4 carbon atoms, are most preferred.

20 The term "alkoxy" refers to an alkyl group bonded through an oxygen (-O-). The term "alkylthio" refers to an alkyl group bonded through a sulfur (-S-).

The term "substituted alkyl" refers to an alkyl chain as defined above having one, two, or three  
25 substituents selected from halo, cyano, hydroxy, alkoxy, alkylthio, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)<sub>2</sub>, carboxy and -CO<sub>2</sub>-alkyl.

The term "alkylene" refers to a bridge of 1 to 10 carbons such as -CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>4</sub>-, etc.

30 The term "substituted alkylene" refers to an alkylene bridge as previously defined having one, two, or

three substituents selected from alkyl, substituted alkyl, halo, cyano, hydroxy, alkoxy, alkylthio, amino -NH(alkyl), -NH(cycloalkyl), -N(alkyl)<sub>2</sub>, carboxy, -CO<sub>2</sub>-alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclo, and heteroaryl.



The term "alkenyl" refers to a bridge of 2 to 10 carbons containing at least one double bond such as -CH=CH-, -CH<sub>2</sub>-CH=CH-, etc.


The term "substituted alkenyl" refers to a bridge of 2 to 10 carbons containing at least one double bond as defined previously having, one, two or three substituents selected from alkyl, substituted alkyl, halo, cyano, hydroxy, alkoxy, alkylthio, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)<sub>2</sub>, carboxy, -CO<sub>2</sub>-alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclo, and heteroaryl.

the term "alkynyl" refers to a bridge of 2 to 10 carbons containing at least one triple bond such as -C≡C-, -CH<sub>2</sub>-C≡C-, etc.

The term "substituted alkynyl" refers to a bridge of 2 to 10 carbons containing at least one triple bond as defined previously having one, two, or three substituents selected from alkyl, substituted alkyl, halo, cyano, hydroxy, alkoxy, alkylthio, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)<sub>2</sub>, carboxy, -CO<sub>2</sub>-alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclo and heteroaryl.

The term "cycloalkyl" refers to cyclic hydrocarbon groups of 3 to 9 carbon atoms, preferably 3 to 7 carbon atoms, which can be fully saturated or partially unsaturated. Also included within this definition are bicyclic cycloalkyl rings having a fused phenyl ring such


as  or  and bicyclic rings having a carbon-carbon bridge of 3 or 4 carbons such as




- 5 The term "substituted cycloalkyl" refers to such cycloalkyl groups as defined above having one, two or three substituents selected from alkyl, substituted alkyl, halo, cyano, hydroxy, alkoxy, alkylthio, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)<sub>2</sub>, carboxy, -CO<sub>2</sub>-alkyl, keto, =N-OH, =N-O-lower alkyl, and a five or six membered ketal, i.e. 1,3-dioxolane or 1,3-dioxane.

The term "aryl" refers to phenyl, 1-naphthyl or 2-naphthyl, with phenyl being preferred.

- 15 The term "substituted aryl" refers to such aryl groups as defined previously having one, two or three substituents selected from alkyl, substituted alkyl, halo, cyano, hydroxy, alkoxy, alkylthio, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)<sub>2</sub>, carboxy, -CO<sub>2</sub>-alkyl, keto, -(C=O)NH<sub>2</sub>, -(C=O)NH(alkyl), -(C=O)NH(cycloalkyl), -(C=O)N(alkyl)<sub>2</sub>, -(C=O)alkyl, -SO<sub>2</sub>NH<sub>2</sub>, -O-(C=O)alkyl, -NH<sub>2</sub>-CH<sub>2</sub>-carboxy, -NH-CH<sub>2</sub>-CO<sub>2</sub>-alkyl, and a five or six membered ring containing two

oxygen atoms such as  or




. Substituted phenyl having one, two or

three substituents as described above is the preferred substituted aryl.

The term "halo" refers to chloro, bromo, fluoro and iodo.

- 5       The term "heterocyclo" refers to substituted and unsubstituted fully saturated or partially unsaturated 3 to 7 membered monocyclic groups, 7 to 11 membered bicyclic groups and 10 to 15 membered tricyclic groups which have at least one heteroatom (O, S or N) in at least one of the rings. Each ring of the heterocyclo group containing a heteroatom can contain one or two oxygen or sulfur atoms and/or from one to four nitrogen atoms provided that the total number of heteroatoms in each ring is four or less, and further provided that each ring contains at least one carbon atom. The fused ring completing the bicyclic and tricyclic groups may be a cycloalkyl, substituted cycloalkyl, aryl or substituted aryl as defined above. The bicyclic ring may also be formed by having a bridge of 2 or 3 carbons between available carbon atoms or between an available carbon and



- nitrogen atoms such as       The nitrogen and sulfur atoms may optionally be oxidized and the nitrogen atom may optionally be quaternized. The heterocyclo group may be attached at any available nitrogen or carbon atom. The heterocyclo ring may contain one, two or three substituents attached to an available carbon or nitrogen atom selected from alkyl, substituted alkyl, halo, cyano, hydroxy, alkoxy, alkylthio, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)<sub>2</sub>, carboxy, -CO<sub>2</sub>-alkyl, keto, -(C=O)NH<sub>2</sub>, -(C=O)NH(alkyl), -(C=O)NH(cycloalkyl), -(C=O)N(alkyl)<sub>2</sub>, -(C=O)alkyl, -O-(C=O)alkyl, -NH-CH<sub>2</sub>-carboxy,

-NH-CH<sub>2</sub>-CO<sub>2</sub>-alkyl, =N-OH, =N-O-alkyl, and a five or six membered ring, i.e., 1,3-dioxolane or 1,3-dioxane.

Exemplary monocyclic groups include azetidiny, pyrrolidinyl, oxetanyl, imidazolinyl, oxazolidinyl, isoxazolinyl, thiazolidinyl, isothiazolidinyl, tetrahydrofuranyl, piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodinyl, 2-oxoazepinyl, azepinyl, 4-piperidonyl, tetrahydropyranyl, morpholinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, 1,3-dioxolane and tetrahydro-1,1-dioxothienyl and the like. Exemplary bicyclic heterocyclo groups include quinuclidinyl.

The term "heteroaryl" refers to substituted and unsubstituted aromatic 5 or 6 membered monocyclic groups, 9 or 10 membered bicyclic groups, and 11 to 14 membered tricyclic groups which have at least one heteroatom (O, S or N) in at least one of the rings. Each ring of the heteroaryl group containing a heteroatom can contain one or two oxygen or sulfur atoms and/or from one to four nitrogen atoms provided that the total number of heteroatoms in each ring is four or less. The fused rings completing the bicyclic and tricyclic groups may be a cycloalkyl, substituted cycloalkyl, aryl or substituted aryl as defined above. The nitrogen and sulfur atoms may optionally be oxidized and the nitrogen atoms may optionally be quaternized. Heteroaryl groups which are bicyclic or tricyclic must include at least one fully aromatic ring but the other fused ring or rings may be aromatic or non-aromatic. The heteroaryl group may be attached at any available nitrogen or carbon atom of any ring. The heteroaryl ring system may contain one, two or three substituents attached to an available carbon or nitrogen atom selected from alkyl, substituted alkyl, halo, cyano, hydroxy, alkoxy, alkylthio, amino,

-NH(alkyl), -NH(cycloalkyl), -N(alkyl)<sub>2</sub>, carboxy, -CO<sub>2</sub>-alkyl, keto, -(C=O)NH<sub>2</sub>, -(C=O)NH(alkyl), -(C=O)NH(cycloalkyl), -(C=O)N(alkyl)<sub>2</sub>, -(C=O)alkyl, -O-(C=O)alkyl, -NH-CH<sub>2</sub>-carboxy, -NH-CH<sub>2</sub>-CH<sub>2</sub>-CO<sub>2</sub>-alkyl, and a five or six membered ring, i.e., 1,3-dioxolane or 1,3-dioxane.

Exemplary monocyclic heteroaryl groups include pyrrolyl, pyrazolyl, pyrazolinyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, isothiazolyl, furanyl, thienyl, oxadiazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, and the like.

Exemplary bicyclic heteroaryl groups include indolyl, benzothiazolyl, benzoxazolyl, benzothienyl, quinolinyl, tetrahydroisoquinolinyl, isoquinolinyl, benzimidazolyl, benzopyranyl, indolizinyl, benzofuranyl, chromonyl, coumarinyl, benzopyranyl, cinnolinyl, quinoxalinyl, indazolyl, pyrrolopyridyl, furopyridinyl, dihydroisoindolyl, tetrahydroquinolinyl and the like.

Exemplary tricyclic heteroaryl groups include carbazolyl, benzidolyl, phenanthrolinyl, acridinyl, phenanthridinyl, xanthenyl and the like.

Throughout the specification, groups and substituents thereof may be chosen to provide stable moieties and compounds.

The compounds of formulas I, II and III form salts which are also within the scope of this invention. Reference to a compound of formulas I, II or III herein is understood to include reference to salts thereof, unless otherwise indicated. The term "salt(s)", as employed herein, denotes acidic and/or basic salts formed with inorganic and/or organic acids and bases. In addition, when a compound of formula I contains a both a basic moiety, such as, but not limited to a pyridine or imidazole, and an acidic moiety, such as, but not limited to a carboxylic acid, zwitterions ("inner salts") may be



formed and are included within the term "salt(s)" as used herein. Pharmaceutically acceptable (i.e., non-toxic, physiologically acceptable) salts are preferred, although other salts are also useful, e.g., in isolation or purification steps which may be employed during preparation. Salts of the compounds of the formulas I, II and III may be formed, for example, by reacting a compound I, II or III with an amount of acid or base, such as an equivalent amount, in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

The compounds of formulas I, II and III which contain a basic moiety, such as, but not limited to an amine or a pyridine or imidazole ring, may form salts with a variety of organic and inorganic acids. Exemplary acid addition salts include acetates (such as those formed with acetic acid or trihaloacetic acid, for example, trifluoroacetic acid), adipates, alginates, ascorbates, aspartates, benzoates, benzenesulfonates, bisulfates, borates, butyrates, citrates, camphorates, camphorsulfonates, cyclopentanepropionates, digluconates, dodecylsulfates, ethanesulfonates, fumarates, glucoheptanoates, glycerophosphates, hemisulfates, heptanoates, hexanoates, hydrochlorides (formed with hydrochloric acid), hydrobromides (formed with hydrogen bromide), hydroiodides, 2-hydroxyethanesulfonates, lactates, maleates (formed with maleic acid), methanesulfonates (formed with methanesulfonic acid), 2-naphthalenesulfonates, nicotines, nitrates, oxalates, pectinates, persulfates, 3-phenylpropionates, phosphates, picrates, pivalates, propionates, salicylates, succinates, sulfates (such as those formed with sulfuric acid), sulfonates (such as those mentioned herein), tartrates, thiocyanates, toluenesulfonates such as tosylates, undecanoates, and the like.

The compounds of formulas I, II and III which contain an acidic moiety, such as, but not limited to a carboxylic acid, may form salts with a variety of organic and inorganic bases. Exemplary basic salts include

5 ammonium salts, alkali metal salts such as sodium, lithium, and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases (for example, organic amines) such as benzathines, dicyclohexylamines, hydrabamines (formed with N,N-

10 bis(dehydroabietyl)ethylenediamine), N-methyl-D-glucamines, N-methyl-D-glucamides, t-butyl amines, and salts with amino acids such as arginine, lysine and the like. Basic nitrogen-containing groups may be

15 quaternized with agents such as lower alkyl halides (e.g. methyl, ethyl, propyl, and butyl chlorides, bromides and iodides), dialkyl sulfates (e.g. dimethyl, diethyl, dibutyl, and diamyl sulfates), long chain halides (e.g. decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides), aralkyl halides (e.g. benzyl and phenethyl

20 bromides), and others.

Prodrugs and solvates of the compounds of the invention are also contemplated herein. The term "prodrug", as employed herein, denotes a compound which, upon administration to a subject, undergoes chemical

25 conversion by metabolic or chemical processes to yield a compound of formulas I, II or III or a salt and/or solvate thereof. Solvates of the compounds of formulas I, II and III are preferably hydrates.

Compounds of the formulas I, II and III and salts

30 thereof, may exist in their tautomeric form (for example, as an amide or imino ether). All such tautomeric forms are contemplated herein as part of the present invention.

All stereoisomers of the present compounds, such as those which may exist due to asymmetric carbons on

35 various substituents, including enantiomeric forms (which

may exist even in the absence of asymmetric carbons) and diastereomeric forms, are contemplated within the scope of this invention. Individual stereoisomers of the compounds of the invention may, for example, be

5 substantially free of other isomers, or may be admixed, for example, as racemates or with all other, or other selected, stereoisomers. The chiral centers of the present invention can have the S or R configuration as defined by the IUPAC 1974 Recommendations.

10

#### Methods of Preparation

The compounds of the present invention may be prepared by methods such as those illustrated in the following Scheme I to IV. Solvents, temperatures,


15 pressures, and other reaction conditions may readily be selected by one of ordinary skill in the art. Starting materials are commercially available or readily prepared by one of ordinary skill in the art.

High Speed Analoging (HSA) may be employed in the

20 preparation of compounds, for example, where the intermediates possess a carboxylic acid group or activated aromatic position, such as the 4 position of a 4-halopyridine. In the same manner, substitutions on the fused five membered ring, such as pyrazoles, imidazoles,

25 and triazoles, may also be achieved through HSA.

## SCHEME I



- 5           Compounds of formulas Ia, IIa and IIIa [wherein E is E<sub>1</sub> or E<sub>2</sub>, X is X<sub>1</sub>, X<sub>2</sub> or X<sub>3</sub>, Y is nitrogen, and Z is C(R<sub>7</sub>)] can be prepared via the aminolysis or
- 10           esterification of a compound of formula IV using an appropriate carboxylic acid activating reagent and an appropriate amine or alcohol of the formulas X<sub>1</sub>-H, X<sub>2</sub>-H or X<sub>3</sub>-H in an inert solvent. Exemplary carboxylic acid
- 15           activating agents include carbonyldiimidazole, dicyclohexylcarbodiimide, pentofluorophenol trifluoroacetate, or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide. Exemplary inert solvents include
- 20           ethers, including tetrahydrofuran and dioxane, N,N-dimethylformamide, acetonitrile, or methylene chloride.
- Compounds of formula IV can be prepared by the hydrolysis of compounds of formula V using a hydroxide
- source. Exemplary hydroxide sources include sodium

hydroxide or lithium hydroxide. Exemplary solvents include water, alcohols, and mixtures of ethers/water.

Compounds of formula V can be prepared by reacting compounds of formula VI with an amine, thiol or alcohol of the formulas E<sub>1</sub>-H or E<sub>2</sub>-H. The reaction may be performed in an inert solvent as appropriate, such as ethanol or N,N-dimethylformamide, in the presence of an appropriate base, such as triethylamine for amines and sodium hydride for thiol or alcohols, and is typically performed at elevated temperatures.


Compounds of formula VI can be prepared from compounds of either formula VIII or formula VII by reacting with an appropriate dehydrating agent typically under elevated temperatures. Exemplary dehydrating agents include POCl<sub>3</sub>, PCl<sub>5</sub>, SOCl<sub>2</sub> and oxalyl chloride.

Compounds of formula VII can be prepared from compounds of formula VIII via an intramolecular cyclization typically performed at elevated temperatures in an inert solvent as appropriate or in neat form.

Compounds of formula VIII can be prepared by combining compounds of formula X and IX either neat or in an inert solvent as appropriate, typically such reaction is performed at elevated temperatures.

Compounds of formula X and formula IX are either commercially available or available via methods known to one skilled in the art. For example, compounds of formula X may be prepared as described in French Patent 1,403,372 [Chemical Abstracts, 1965, Volume 63, 14871a].

## SCHEME II



5

Compounds of formula Ib, IIb or IIIb [wherein E is E<sub>1</sub> or E<sub>2</sub>, X is X<sub>1</sub>, X<sub>2</sub> or X<sub>3</sub>, and Y and Z are both nitrogen] can be prepared via the aminolysis or esterification of a compound of formula XI using an appropriate carboxylic acid activating reagent and an appropriate amine or alcohol of the formula X-H in an inert solvent.

Exemplary carboxylic acid activating agents include carbonyldiimidazole, dicyclohexylcarbodiimide, pentofluorophenol trifluoroacetate, or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide. Exemplary inert solvents include ethers, including tetrahydrofuran and dioxane, N,N-dimethylformamide, acetonitrile, or methylene chloride.

Compounds of formula XI can be prepared from by the hydrolysis of compounds of formula XII using a hydroxide source. Exemplary hydroxide sources include sodium hydroxide or lithium hydroxide. Exemplary solvents include water, alcohols, and mixtures of ethers/water.

Compounds of formula XII can be prepared by treating a compound of formula XIII with a diazotizing reagent in an acidic aqueous medium. Sodium nitrite is an exemplary

diazotizing reagent and dilute (1N) HCl is an exemplary reaction solvent.


Compounds of formula XIII can be prepared via the reduction of a compound of formula XIV in an inert solvent. This reduction may, for example, be mediated via a platinum or palladium-catalyzed hydrogenation using platinum or palladium on carbon, hydrogen and an inert solvent such as ethanol or methanol or, alternatively, by use of a stoichiometric reducing agent, such as stannous(II) chloride, in an inert solvent such as ethyl acetate.

Compounds of formula XIV can be prepared by reacting compounds of formula XV with amines of the formula  $R_3NH_2$ . The reaction may be performed in an inert solvent as appropriate, such as ethanol, in the presence of appropriate base, such as triethylamine, and typically under elevated temperatures.

Compounds of formula XV can be prepared by reacting compounds of formula XVI with an amine, thiol or alcohol of the formula E-H. The reaction may be performed in an inert solvent as appropriate, such as N,N-dimethylformamide, in the presence of an appropriate base, such as triethylamine for amines and sodium hydride for thiols or alcohols.

Methods of synthesis of compounds of formulas XVII, XVI, XV, XIV, XIII, XII, and XI are known to one skilled in the art. For example such methodology can be found in US 4,070,362, US 4,003,908, and US 4,048,182. Compounds of formula XVII are either commercially available or prepared by methods known to one skilled in the art.

## Scheme III



5           Compounds of formula Ic, IIc and IIIc [wherein E is E<sub>1</sub> or E<sub>2</sub> and X is X<sub>1</sub>, X<sub>2</sub> or X<sub>3</sub>, Y is C(R<sub>6</sub>) and Z is nitrogen] can be prepared via the aminolysis or esterification of a compound of formula XVIII using an appropriate carboxylic acid activating reagent and an appropriate amine or alcohol of the formula X-H in an inert solvent. Exemplary carboxylic acid activating agents include carbonyldiimidazole, dicyclohexylcarbodiimide, pentofluorophenol trifluoroacetate, or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide. Exemplary inert solvents include ethers, including tetrahydrofuran and dioxane, N,N-dimethylformamide, acetonitrile, or methylene chloride.

10           

15           

20           Compounds of formula XVIII can be prepared from by the hydrolysis of compounds of formula XIX using a hydroxide source. Exemplary hydroxide sources include sodium hydroxide or lithium hydroxide. Exemplary solvents include water, alcohols, and mixtures of ethers/water.


25           Compounds of formula XIX can be prepared from the condensation of compounds of formula XIII with an activated ester derivative from an acid of the formula



$R_6\text{-CO}_2\text{H}$  under basic conditions in an inert solvent typically under elevated temperatures. Exemplary activated esters include acid chlorides derived from  $R_6\text{-CO}_2\text{H}$ , N,N-dialkylamide acetals (including, for example, N,N-dimethylformamide dimethyl acetal) and activated esters derived from the reaction of  $R_6\text{-CO}_2\text{H}$  with exemplary carboxylic acid activating agents such as carbonyldiimidazole, dicyclohexylcarbodiimide, pentofluorophenol trifluoroacetate, or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide. Exemplary bases include sodium hydride, potassium hydride, cesium carbonate, potassium carbonate, potassium hexamethyldisilazide, and potassium *t*-butoxide. Exemplary inert solvents include ethers, N,N-dimethylformamide, and acetonitrile.

Compounds of formula XIII are prepared as discussed in Scheme II.

Scheme IV



Compounds of formula Ia, IIa and IIIa [wherein E is  $E_1$  or  $E_2$ , X is  $X_1$ ,  $X_2$  or  $X_3$ , Y is nitrogen, and Z is  $C(R_7)$ ] can also be prepared by reacting compounds of formula XX

with an amine, thiol or alcohol of the formula E-H. The reaction may be performed in an inert solvent as appropriate, such as N,N-dimethylformamide, in the presence of an appropriate base, such as triethylamine  
5 for amines and sodium hydride for alcohols, and typically under elevated temperatures..

Compounds XX can be prepared by reacting compounds of formula XXI with an appropriate amine or alcohol of the formula X-H in an inert solvent in the presence of an  
10 appropriate base such as triethylamine. Exemplary inert solvents include ethers, including tetrahydrofuran and dioxane, N,N-dimethylformamide, acetonitrile, or methylene chloride.

Compounds XXI can be prepared from compounds of  
15 formula XXII by reacting with an appropriate dehydrating agent typically under elevated temperatures. Exemplary dehydrating agents include POCl<sub>3</sub>, PCl<sub>5</sub>, SOCl<sub>2</sub> and oxalyl chloride.

Compounds XXII can be prepared from by the  
20 hydrolysis of compounds of formula VII using a hydroxide source. Exemplary hydroxide sources include sodium hydroxide or lithium hydroxide. Exemplary solvents include water, alcohols, and mixtures of ethers/water.

Compound VII can be prepared as described in Scheme  
25 I.

#### Preferred Compounds and Method

The following compounds of formula I and II are  
30 preferred:

Y is nitrogen.





Z is nitrogen or C(R<sub>7</sub>).

E<sub>1</sub> is -O-R<sub>1</sub>, -NH-A<sub>1</sub>-cycloalkyl, -NH-A<sub>1</sub>-heterocyclo,  
or -NH-A<sub>1</sub>-heteroaryl.

$E_2$  is  $-NH-A_1$ -alkoxy,  $-NH-A_1-N \begin{matrix} R_{15} \\ R_{16} \end{matrix}$ ,  $-NH-A_1$ -substituted phenyl, or  $-NH-A_1-CO_2$ -alkyl.

$R_1$  is  $-A_1$ -substituted phenyl.

$X_1$  is  $-O-A_1$ -heteroaryl,  $-O-A_1$ -heterocyclo,



10  $R_7$  is hydrogen.

$R_4$  is hydrogen.

$R_3$  is straight or branched chain alkyl of 1 to 4 carbons.

15  $R_5$  is hydrogen, alkyl,  $-CO_2$ -alkyl,  $-A_1$ -phenyl, or  $-A_1$ -heteroaryl wherein alkyl is straight or branched chain of 1 to 4 carbons.

20  $R_2$  is  $-(C=O)R_9$ ,  $-S-R_9$ ,  $-O-R_9$ ,  $-N(R_9)(R_{10})$ ,  $-NR_{11}CO_2R_{19}$ ,  $-(C=O)N(R_{11})CH_2CO_2R_{19}$ ,  $-NR_{11}(C=O)R_{19}$ ,  $-CO_2R_9$ , nitrogen when  $A_2$  is alkynyl ending in a triple bond,  $-(C=O)N(R_{12})(R_{13})$ , phenyl, substituted phenyl, cycloalkyl, heterocyclo, heteroaryl, cycloalkyl- $A_3$ -phenyl, phenyl- $A_3$ -heteroaryl, heterocyclo- $A_3$ -phenyl, or heterocyclo- $A_3$ -heterocyclo.

$R_{25}$  is  $-S-R_9$ ,  $-NR_{11}CO_2R_{19}$ , nitrogen when  $A_2$  is alkynyl ending in a triple bond,  $-(C=O)N(R_{11})CH_2CO_2R_{19}$ ,

-NR<sub>11</sub>(C=O)R<sub>19</sub>, -CO<sub>2</sub>R<sub>9</sub>, -(C=O)N(R<sub>12</sub>)(R<sub>13</sub>), phenyl, substituted phenyl, cycloalkyl, heterocyclo, heteroaryl, cycloalkyl-A<sub>3</sub>-phenyl, heterocyclo-A<sub>3</sub>-phenyl, phenyl-A<sub>3</sub>-heteroaryl or heterocyclo-A<sub>3</sub>-heterocyclo.

5        A<sub>1</sub> is an alkylene or substituted alkylene bridge of 1 to 6 carbons wherein said substituent is a straight or branched chain alkyl of 1 to 4 carbons.

         A<sub>2</sub> is a direct bond, an alkylene or substituted  
10        alkylene bridge of 1 to 6 carbons wherein said substituent is one or two members selected from alkyl, phenyl, substituted phenyl, -CO<sub>2</sub>-alkyl, carboxy, hydroxy, -NH-(C=O)-alkyl, and -CH<sub>2</sub>-(C=O)-NH<sub>2</sub>, an alkenyl bridge of 2 to 4 carbons having one double bond, or an alkynyl  
15        bridge of 2 to 3 carbons having one triple bond wherein alkyl is straight or branched chain of 1 to 4 carbons.


         The term "heterocyclo" in the preferred definitions refers to a substituted or unsubstituted fully saturated or partially saturated 5 to 7 membered monocyclic rings containing one or two heteroatoms selected from oxygen,  
20        sulfur and nitrogen and bicyclic rings wherein the monocyclic ring as defined above is fused to a phenyl or substituted phenyl or wherein a bridge of 2 or 3 carbons is present between available carbon and nitrogen atom. The nitrogen and sulfur atoms may optionally be oxidized  
25        and the nitrogen atom may optionally be quaternized. The heterocyclo group may be attached at any available nitrogen or carbon atom. The heterocyclo ring may contain one or two substituents attached to an available carbon or nitrogen atom selected from alkyl, keto and  
30        -CO<sub>2</sub>-alkyl, wherein alkyl is straight or branched chain of 1 to 4 carbons.

         The term "heteroaryl" in the preferred definitions refers to a substituted or unsubstituted aromatic 5 or 6  
35        membered monocyclic ring containing one or two oxygen or sulfur atoms and/or from one to four nitrogen atoms

provided that the total number of heteroatoms in the ring is four or less, and bicyclic rings wherein the monocyclic ring as defined above is fused to a phenyl or substituted phenyl. The nitrogen and sulfur atoms may  
 5 optionally be oxidized and the nitrogen atom may optionally be quaternized. The heteroaryl group may be attached at any available nitrogen or carbon atom. The heteroaryl ring may contain one or two substituents  
 10 attached to an available carbon or nitrogen atom selected from straight or branched chain alkyl of 1 to 4 carbons and halo.

The term "cycloalkyl" in the preferred definitions refers to a fully saturated cyclic hydrocarbon group of 3 to 7 carbons and such cycloalkyl rings fused to a phenyl  
 15 ring or such cycloalkyl rings 5 to 7 carbons having a carbon-carbon bridge of 3 or 4 carbons.


The term "substituted phenyl" refers to a phenyl ring having one, two, or three substituents selected from alkyl, halo, hydroxy, trifluoromethyl, alkoxy of 1 to 4  
 20 carbons, -N(alkyl)<sub>2</sub>, and SO<sub>2</sub>NH<sub>2</sub> wherein alkyl is straight or branched chain of 1 to 4 carbons, and a phenyl ring substituted with a fused five membered ketal, i.e.



25 R<sub>9</sub>, R<sub>10</sub>, R<sub>11</sub>, R<sub>12</sub>, R<sub>13</sub>, R<sub>15</sub>, R<sub>16</sub> and R<sub>19</sub> are independently selected from the group consisting of hydrogen, straight or branched chain alkyl of 1 to 4 carbons, phenyl, substituted phenyl and -A<sub>1</sub>-phenyl.

A<sub>3</sub> is a direct bond, an alkylene bridge of 1 to 6

30 carbons, or  $\text{—CH}_2\text{—C(=O)—}$ .



is a 5 to 7 membered heterocyclo ring which can contain an additional nitrogen atom or can contain an oxygen or sulfur atom.


- 5  $R_{21}$  is attached to an available carbon or nitrogen atom and is hydrogen, straight or branched chain alkyl of 1 to 4 carbons, hydroxy or amino.

- 10  $R_{22}$  is attached to an available carbon or nitrogen atom and is keto,  $-(C=O)R_{23}$ ,  $-CO_2R_{23}$ ,  $-NH-(C=O)-R_{23}$ ,  $-N(alkyl)_2$ ,  $-A_1$ -hydroxy,  $-A_1-N(R_9)(R_{10})$ ,  $-A_1$ -alkoxy,  $-A_2$ -phenyl,  $-A_2$ -substituted phenyl, or  $-A_2$ -heteroaryl wherein alkyl is straight or branched chain of 1 to 4 carbons and alkoxy is such an alkyl bonded through an oxygen.


$n$  is one or two.

$m$  is zero or one.

- 15  $R_{23}$  is alkyl,  $-N(R_9)(R_{10})$ , or  $-A_2$ -heteroaryl wherein alkyl is straight or branched chain of 1 to 4 carbons.





represents a fused bicyclic ring wherein

the monocyclic ring  is defined previously and






represents a substituted phenyl having two carbon

- 20 atoms in common with the monocyclic ring .




represents a spiro ring wherein the

monocyclic ring  is defined previously and  represents a heterocyclo ring having a common carbon with

the monocyclic ring .

Also preferred is the method of treating a cGMP-associated condition, particularly erectile dysfunction, with the preferred compounds of formulas I and II as defined above or with a compound of formula III wherein:

5 Y, Z, E<sub>2</sub>, R<sub>4</sub> and R<sub>3</sub> are as defined for the preferred compounds of formula II, and



10

R<sub>5</sub>, A<sub>1</sub>, A<sub>2</sub>, R<sub>9</sub>, R<sub>10</sub>, and  $\begin{array}{c} \text{---N} \\ | \\ \text{(R}_{21}\text{)}_n \end{array}$  are as defined for the preferred compounds of formula II.

### More Preferred Compounds And Method

15

The following compounds of formula I and II are more preferred:


Y is nitrogen.

Z is CH.

20

R<sub>3</sub> is ethyl.


R<sub>4</sub> is hydrogen.



25


E<sub>2</sub> is -NH-CH<sub>2</sub>-disubstituted phenyl.

The term "disubstituted phenyl" refers to a phenyl ring having two substituents independently selected from halogen and methoxy or wherein said disubstituted phenyl




5

$X_1$  is  $-O-A_1$ -heterocyclo,  $-O-A_1$ -heteroaryl,  $-NH-A_2-R_2$ ,



$X_2$  is  $-O-A_1$ -heterocyclo,  $-O-A_1$ -heteroaryl,  $-NH-A_2-R_{25}$ ,



10

$A_1$ ,  $A_2$ ,  $R_2$ ,  $R_{25}$ , "heterocyclo" and "heteroaryl" are as defined in the preferred definitions.

Also more preferred is the method of treating a cGMP associated condition, particularly erectile dysfunction, with the more preferred compounds of formula I and II as defined above or with a compound of formula III wherein:

$Y$ ,  $Z$ ,  $E_2$ ,  $R_4$  and  $R_3$  are as defined for the preferred compounds of formula II, and



$X_3$  is  $-\text{NH}-\text{A}_2-\text{OR}_9$  wherein  $\text{A}_2$  and  $\text{R}_9$  are as defined for the preferred compounds of formula II.

5

### Most Preferred Compounds And Method

The following compounds of formulas I and II are most preferred:



Y is nitrogen.

10

Z is CH.


$\text{R}_3$  is ethyl.


$\text{R}_4$  is hydrogen.



$\text{X}_1$  and  $\text{X}_2$  are independently selected from the group consisting of


20





Also most preferred is the method of treating a cGMP associated condition, particularly erectile dysfunction, with the most preferred compounds of formulas I and II as defined above or with a compound of formula III wherein:

- 5        Y, Z, E<sub>2</sub>, R<sub>4</sub> and R<sub>3</sub> are as defined the most preferred compounds of formula II, and



10

### Utility

- The compounds of the present invention inhibit cGMP PDE, and in particular are potent and selective inhibitors of cGMP PDE V. The present compounds are
- 15        useful in the treatment of cGMP-associated conditions. A "cGMP-associated condition", as used herein, denotes a disorder which can be treated by inhibiting cGMP PDE or elevating the level of cGMP in a subject, wherein treatment comprises prevention, partial alleviation or
- 20        cure of the disorder. Inhibition of cGMP PDE or elevation of the cGMP level may occur locally, for example, within certain tissues of the subject, or more extensively throughout a subject being treated for such a disorder. Treatment may be facilitated wherein elevation
- 25        of the cGMP level potentiates additional beneficial therapeutic effects, such as where elevation of the cGMP level potentiates the effects of endothelium-derived relaxing factor.

- The compounds of the present invention are useful
- 30        for the treatment of a variety of cardiovascular diseases including, but not limited to, hypertension, angina (stable, unstable, and variant), (congestive) heart failure, restenosis, atherosclerosis, and dyslipidemia, as well as reduced blood vessel patency, thrombus, both

venous and arterial, myocardial infarction, peripheral vascular disease, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma, diseases characterized by disorders of gut motility, and forms of cancer responsive to the inhibition of cGMP PDE. In addition, these compounds are useful in the treatment of sexual dysfunction in both men (erectile dysfunction, for example, due to diabetes mellitus, spinal cord injury, radical prostatectomy, psychogenic etiology or any other cause) and women by improving blood flow to the genitalia, especially, the corpus cavernosum.

The present invention thus provides methods for the treatment of cGMP-associated conditions, comprising the step of administering to a subject in need thereof at least one compound of formulas I, II or III in an amount effective therefor. Other therapeutic agents such as those described below may be employed with the inventive compounds in the present methods. In the methods of the present invention, such other therapeutic agent(s) may be administered prior to, simultaneously with or following the administration of the compound(s) of the present invention.

The present invention also provides pharmaceutical compositions comprising at least one of the compounds of formulas I, II or III capable of treating a cGMP-associated condition in an amount effective therefor, and a pharmaceutically acceptable vehicle or diluent. The compositions of the present invention may contain other therapeutic agents as described below, and may be formulated, for example, by employing conventional solid or liquid vehicles or diluents, as well as pharmaceutical additives of a type appropriate to the mode of desired administration (for example, excipients, binders, preservatives, stabilizers, flavors, etc.) according to

techniques such as those well known in the art of pharmaceutical formulation.

The compounds of formulas I, II and III may be administered by any suitable means, for example, orally, such as in the form of tablets, capsules, granules or powders; sublingually; buccally; parenterally, such as by subcutaneous, intravenous, intramuscular, or intrasternal injection or infusion techniques (e.g., as sterile injectable aqueous or non-aqueous solutions or suspensions); nasally such as by inhalation spray; topically, such as in the form of a cream or ointment; or rectally such as in the form of suppositories; in dosage unit formulations containing non-toxic, pharmaceutically acceptable vehicles or diluents. The present compounds may, for example, be administered in a form suitable for immediate release or extended release. Immediate release or extended release may be achieved by the use of suitable pharmaceutical compositions comprising the present compounds, or, particularly in the case of extended release, by the use of devices such as subcutaneous implants or osmotic pumps. The present compounds may also be administered liposomally.

Exemplary compositions for oral administration include suspensions which may contain, for example, microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners or flavoring agents such as those known in the art; and immediate release tablets which may contain, for example, microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate and/or lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants such as those known in the art. The compounds of formulas I, II and III may also be delivered through the oral cavity by sublingual and/or buccal

administration. Molded tablets, compressed tablets or freeze-dried tablets are exemplary forms which may be used. Exemplary compositions include those formulating the present compound(s) with fast dissolving diluents  
5 such as mannitol, lactose, sucrose and/or cyclodextrins. Also included in such formulations may be high molecular weight excipients such as celluloses (Avicel®) or polyethylene glycols (PEG). Such formulations may also include an excipient to aid mucosal adhesion such as  
10 hydroxy propyl cellulose (HPC), hydroxy propyl methyl cellulose (HPMC), sodium carboxy methyl cellulose (SCMC), maleic anhydride copolymer (e.g., Gantrez®), and agents to control release such as polyacrylic copolymer (e.g., Carbopol® 934). Lubricants, glidants, flavors, coloring  
15 agents and stabilizers may also be added for ease of fabrication and use.

Exemplary compositions for nasal aerosol or inhalation administration include solutions in saline which may contain, for example, benzyl alcohol or other  
20 suitable preservatives, absorption promoters to enhance bioavailability, and/or other solubilizing or dispersing agents such as those known in the art.

Exemplary compositions for parenteral administration include injectable solutions or  
25 suspensions which may contain, for example, suitable non-toxic, parenterally acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution, an isotonic sodium chloride solution, or other suitable dispersing or wetting and suspending agents, including  
30 synthetic mono- or diglycerides, and fatty acids, including oleic acid.

Exemplary compositions for rectal administration include suppositories which may contain, for example, a  
35 suitable non-irritating excipient, such as cocoa butter, synthetic glyceride esters or polyethylene glycols, which

are solid at ordinary temperatures, but liquefy and/or dissolve in the rectal cavity to release the drug.

Exemplary compositions for topical administration include a topical carrier such as Plastibase (mineral oil gelled with polyethylene).

The effective amount of a compound of the present invention may be determined by one of ordinary skill in the art, and includes exemplary dosage amounts for an adult human of from about 0.05 to 100 mg/kg of body weight of active compound per day, which may be administered in a single dose or in the form of individual divided doses, such as from 1 to 4 times per day. It will be understood that the specific dose level and frequency of dosage for any particular subject may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the species, age, body weight, general health, sex and diet of the subject, the mode and time of administration, rate of excretion, drug combination, and severity of the particular condition. Preferred subjects for treatment include animals, most preferably mammalian species such as humans, and domestic animals such as dogs, cats, horses and the like, subject to cGMP-associated conditions.

The compounds of the present invention may be employed alone or in combination with each other and/or other suitable therapeutic agents useful in the treatment of cGMP-associated conditions such as other cGMP PDE inhibitors, particularly other cGMP PDE V inhibitors, prostanoids,  $\alpha$ -adrenergic agonists, endothelin antagonists, angiotensin II (especially, subtype AT<sub>1</sub>) antagonists, angiotensin converting enzyme (ACE) inhibitors, renin inhibitors, and serotonin (5-HT<sub>2c</sub>) agonists.

Exemplary such other therapeutic agents include the following: phentolamine, yohimbine, papaverine, apomorphine, sildenafil (see *Drugs of the Future*, 22, 138-143 (1997)), pyrazolopyrimidinones as described in  
5 U.S. Patent Nos. 5,272,147; 5,250,534; 5,426,107; and 5,346,901, quinazolinones as described in U.S. Patent No. 5,482,941; AT<sub>1</sub> antagonists selected from losartan, irbesartan, valsartan and candesartan; ET<sub>A</sub> antagonists selected from bosentan, ABT-627, and those described in  
10 U.S. Patent No. 5,612,359 and U.S. Patent Application Serial No. 60/035,832, filed January 30, 1997; PDE V inhibitors selected from imidazoquinazolines (see WO 98/08848), carbazoles (see WO 97/03675, WO 97/03985 and WO 95/19978), imidazopurinones (see WO 97/19947),  
15 benzimidazoles (see WO 97/24334), pyrazoloquinolines (see U.S. Patent No. 5,488,055), anthranilic acid derivatives (see WO 95/18097), fused heterocycles (see WO 98/07430) and thienopyrimidines (see DE 19632423); and 5-HT<sub>2c</sub> agonists selected from indoles (see *J. Med. Chem.*, 40, 2762-2769 (1997), EP 655440 and EP 657426).  
20

The above other therapeutic agents, when employed in combination with the compounds of the present invention, may be used, for example, in those amounts indicated in the Physicians' Desk Reference (PDR) or as  
25 otherwise determined by one of ordinary skill in the art.

The following assay can be employed in ascertaining the degree of activity of a compound as a cGMP PDE inhibitor. Compounds described in the following  
30 Examples have been tested in this assay, and have shown activity.



PDE Scintillation Proximity Assay Protocol

Sonicated human platelet homogenates are prepared by the method of Seiler, et al. (Seiler, S., Gillespie, E., Arnold, A.J., Brassard, C.L., Meanwell, N.A. and Fleming, J.S., "Imidazoquinoline derivatives: potent inhibitors of platelet cAMP phosphodiesterase which elevate cAMP levels and activate protein kinase in platelets," *Thrombosis Research*, **62**: 31-42 (1991)). PDE V is abundant in human platelets, and accounts for approximately 90% of the cGMP hydrolytic activity in the homogenates. When necessary, PDE V can be resolved from other PDE activities in the homogenates by anion exchange chromatography on a fast protein liquid chromatography system (FPLC) using a Mono-Q anion exchange column (Pharmacia) eluted with a linear gradient of 10 mM - 450 mM NaCl.

The phosphodiesterase activity is assayed using a commercially available phosphodiesterase [<sup>3</sup>H]cGMP scintillation proximity (SPA) assay kit (Amersham). The manufacturer's protocol is followed explicitly except that the reactions are carried out at room temperature and 3 mM nonradioactive cGMP is included in the suspension of SPA beads to prevent the synthesis of any additional radioactive products.

25

All documents cited in the present specification are incorporated herein by reference in their entirety.

The following Examples illustrate embodiments of the present invention, and are not intended to limit the scope of the claims.

30


**Abbreviations**

- DMF = dimethylformamide  
DMSO = dimethylsulfoxide
- 5 Et = ethyl  
HPLC = high pressure liquid chromatography  
LRMS = low resolution mass spectrometry  
Me = methyl  
MeOH = methanol
- 10 mp = melting point  
THF = tetrahydrofuran  
tlc = thin layer chromatography  
rt = room temperature  
h = hours
- 15 H<sub>2</sub>O = water  
POCl<sub>3</sub> = phosphorus oxychloride  
EtOH = ethanol  
H<sub>3</sub>PO<sub>4</sub> = phosphoric acid  
HCl = hydrogen chloride
- 20 NaOH = sodium hydroxide  
min = minutes  
Et<sub>3</sub>N = triethylamine  
Et<sub>2</sub>O = ethyl ether  
EtOAc = ethyl acetate
- 25 NaHCO<sub>3</sub> = sodium bicarbonate  
MgSO<sub>4</sub> = magnesium sulfate  
CH<sub>2</sub>Cl<sub>2</sub> = methylene chloride  
Na<sub>2</sub>SO<sub>4</sub> = sodium sulfate  
K<sub>2</sub>CO<sub>3</sub> = potassium carbonate
- 30

### Preparation of Starting Materials

#### Preparation 1


#### Preparation of (1-Ethylpyrazol-5-yl-amino)methylenemalonate diethyl ester




- 5           A neat solution of 5-amino-1-ethylpyrazole (20.0 g, 180 mmol) and diethyl ethoxymethylenemalonate (42.8 g, 198 mmol) was heated at 120 °C for 5 h. This material was used directly without further purification. If needed, the product can be distilled at 154-160 °C (0.1 mm Hg) to afford the title compound as a liquid which solidified to afford the title compound as a pale
- 10   colored solid: mp 50-53 °C.

#### Preparation 2

#### Preparation of 1-Ethyl-4-hydroxy-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid ethyl ester




- 15           (1-Ethylpyrazol-5-yl-amino)methylenemalonate diethyl ester (180 mmol from previous reaction) was dissolved in diphenyl ether (200 mL), and the resulting solution was placed in a preheated oil bath at 255 °C. The reaction solution was heated for 5 h, and then the diphenyl ether was removed via
- 20   distillation. The resulting brown reaction mixture was cooled to room temperature and poured to hexane (1L). Cooling this solution to -78 °C, followed by filtration of the resulting precipitate afforded the title compound as a beige colored needle shaped solid that was >90% pure by HPLC and was used directly (25 g, 60% for 2 steps). A portion was recrystallized using
- 25   ethanol-H<sub>2</sub>O to afford a white solid: mp 85-86 °C; LRMS (m/z) 236 (MH<sup>+</sup>).

Preparation 3Preparation of 4-Chloro-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid ethyl ester

- 5            1-Ethyl-4-hydroxy-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid ethyl ester (15 g, 63.8 mmol) was dissolved in POCl<sub>3</sub> (100 mL), and the resulting solution was heated at reflux for 4 h. The remaining POCl<sub>3</sub> was removed via evaporation under reduced pressure. The residual light brown solid was recrystallized from EtOH-hexane to afford the title compound as a white solid
- 10 (14 g, 55.3 mmol, 87%): HPLC (YMC S5 ODS 4.6x50 mm column, 4 minute gradient- 0% B to 100% B, 4 mL/min flow, solvent A: 10% MeOH-90% H<sub>2</sub>O-0.2% H<sub>3</sub>PO<sub>4</sub>, solvent B: 90% MeOH-10% H<sub>2</sub>O-0.2% H<sub>3</sub>PO<sub>4</sub>) retention time 3.84 minutes showed a purity of 96%; LRMS (m/z) 254 (MH<sup>+</sup>).

15


Preparation 4Preparation of 4-Chloro-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid ethyl ester

- (1-Ethylpyrazol-5-yl-amino)methylenemalonate diethyl ester (10.0 g, 42.6 mmol) was dissolved in 50 mL POCl<sub>3</sub>. This solution was heated at reflux for 10 h before the POCl<sub>3</sub> was removed under reduced pressure. The resulting brown residue was diluted with 5 mL EtOH and extracted with hot hexane (200 mL x 3). The combined organic layers were evaporated under reduced pressure to afford the title compounds which formed light green
- 20
- 25 needle shaped crystals upon standing at room temperature (5.4 g, 21.3

mmol, 50% yield). This material is identical to the one obtained in Preparation 3 ( $^1\text{H}$  NMR,  $^{13}\text{C}$  NMR, MS, and HPLC).


#### Preparation 5

##### 5      (3-Chloro-4-methoxyphenyl)methylamine hydrogen chloride



(4-Methoxyphenyl)methylamine (75.0 g, 0.55 mol) was dissolved in 400 mL diethyl ether. Hydrogen chloride (4.0 M in dioxane, 1.1 mol) was added dropwise with vigorous stirring. After the addition completed, the white solid was filtered and washed thoroughly with diethyl ether. The solid was air dried over night (95.0 g, 100%).


Chlorine gas was bubbled into 400 mL glacial acetic acid with stirring until the weight gained equaled 7% of the starting acetic acid. In a 2 L round bottom flask, 4-methoxybenzylamine hydrogen chloride (32.0 g, 0.18 mol) was suspended in 400 mL glacial acetic acid with vigorous stirring. The chlorine solution (1.5 eq Cl<sub>2</sub>) was added in rapid drops in 30 min at room temperature. The resulted suspension was stirred for another 20 min before N<sub>2</sub> was bubbled in to remove Cl<sub>2</sub> and HCl into a 6 N NaOH trap. The acetic acid was evaporated under reduced pressure to 100 mL. To this white slurry, diethyl ether (300 mL) was used to loosen the solid which was then filtered. The solid was resuspended with 50 mL acetic acid followed by the addition of 50 mL diethyl ether and filtration. This process was repeated twice. The white solid was transferred to a 1 L Erhlenmeyer flask and suspended in 400 mL THF. This suspension was heated to boiling for 10 min before filtration. The undissolved solid was filtered, and twice resuspended in boiling THF (100 mL) with filtration to afford the title compound (27.0 g, 71%) as a white solid. This material contained <2% starting material and <2% dichlorinated material.

Preparation 6General Procedure to Prepare 4-Aminopyrazolopyridine-5-carboxylic acids

The appropriate 4-chloro-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid  
 5 ethyl ester ( 4.0 mmol) and the appropriate amine (4.4 mmol)) were  
 suspended in EtOH (20 mL). To this suspension was added of Et<sub>3</sub>N (2.8 mL,  
 20 mmol), and the resulting solution was heated at reflux for 10 h. To the  
 resulting suspension was added 6 N NaOH (2.7 mL, 16 mmol), and the  
 resulting mixture was heated at reflux for 3 h. EtOH was then removed via  
 10 evaporation under reduced pressure, and the residual white solid was  
 dissolved in 25 mL 0.1 N NaOH. The resulting aqueous mixture was  
 extracted with diethyl ether (3x150 mL), and the organic extracts were  
 discarded. The aqueous layer was acidified with 1 N HCl, and the resulting  
 white precipitate was collected by filtration and washed sequentially with 1 N  
 15 HCl, H<sub>2</sub>O, EtOH and Et<sub>2</sub>O to afford the desired 4-aminopyrazolopyridine.


Using the above described procedure, the following compounds were  
 prepared:

A. 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-  
 20 pyrazolo[3,4-b]pyridine-5-carboxylic acid




4-Chloro-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid ethyl ester and (3-chloro-4-methoxyphenyl)methylamine was used to afford the title compound (87%) as a white solid: mp 250 °C (decomposed).

- 5            B.        (R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid




- 4-Chloro-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid ethyl ester and (R)-2-cyclohexylethylamine was used to afford the title compound  
10        (80%) as a white solid: mp 200 °C (decomposed).

- C. 4-[[[(3,4-Dioxomethylenepheryl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid




- 15            4-Chloro-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid ethyl ester and (3,4-dioxomethylenepheryl)methylamine were used to afford the title compound (80%) as a white solid: mp 230 °C (decomposed).

Preparation 7(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid pentafluorophenyl ester

- 5           The (R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid (0.3 g, 0.95 mmol) was suspended in 2 mL DMF with the subsequent addition of pyridine (0.15 g, 1.9 mmol), pentofluorophenol trifluoroacetate (0.53 g, 1.9 mmol), and a catalytic amount of pentafluorophenol (10 mg). This mixture was stirred at rt for 48 h, and the
- 10           resulting mixture was diluted with EtOAc (50 mL) and washed with 0.1 N HCl (50 mL), 5% NaHCO<sub>3</sub> (3 x 50 mL), and brine (5 x 50 mL). The organic extract was dried over MgSO<sub>4</sub> and concentrated to afford the title compound as a viscous oil (0.44 g, 95%): HPLC (YMC S5 ODS 4.6x50 mm column, 4 minute gradient- 50% B to 100% B, 4.0 mL/min flow, solvent A: 10% MeOH-90%
- 15           H<sub>2</sub>O-0.2% H<sub>3</sub>PO<sub>4</sub>, solvent B: 90% MeOH-10% H<sub>2</sub>O-0.2% H<sub>3</sub>PO<sub>4</sub>) retention time 5.06 minutes showed a purity of 95%; LRMS (m/z) 483 (MH<sup>+</sup>).

The resulting product contained ~5% free pentafluorophenol and was used directly in subsequent reactions.

20

Preparation 8Preparation of 1-Ethyl-4-hydroxy-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid




1-Ethyl-4-hydroxy-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid ethyl ester (10.0 g, 42.6 mmol) was suspended in 50 mL EtOH. To this suspension was added 6 N NaOH (21.3 mL, 3 eq). The resulting light brown solution was heated at reflux for 3 hr before being diluted with 100 mL water. The aqueous solution was extracted with diethyl ether (50 mL x 3) and the organic layer was discarded. The aqueous layer was acidified with 1 N HCl and a white precipitate formed. The title compound was collected by filtration and rinsed with water and diethyl ether affording a white solid (7.9 g, 90%): mp 218 °C (decomposed); LRMS (m/z) 208 (MH<sup>+</sup>).

10


#### Preparation 9

##### 4-Chloro-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid chloride



1-Ethyl-4-hydroxy-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid (0.92 g, 4.4 mmol) was dissolved in 10 mL POCl<sub>3</sub>, and the solution was heated at reflux for 2 hr. The excess POCl<sub>3</sub> was removed under reduced pressure and azeotroped three times with toluene. The brown residue was dissolved in 20 mL CH<sub>2</sub>Cl<sub>2</sub>, Et<sub>3</sub>N (5 eq) and an amine such as 4-aminomethylpyridine (1.2 eq) were then added sequentially. The resulting solution was stirred at rt for 2 h before being diluted with 100 mL water. The organic layer was separated and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (50 mL X 3). The combined organic layer was washed with NaHCO<sub>3</sub> and water. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and the solvent was removed. The residue was chromatographed using silica gel column (5% MeOH-CH<sub>2</sub>Cl<sub>2</sub>) to afford the title compound as a white solid (0.6 g, 44%).

25

Example 1General Procedures for the Preparation of 4-Aminopyrazolopyridines-5-carboxylates and 4-Aminopyrazolopyridines-5-carboxamides

- 5        Method A: To a mixture of the appropriate 4-amino pyrazolopyridine-5-carboxylic acid (5.6 mmol), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (EDAC•HCl, 1.6 g, 8.4 mmol), and 1-hydroxybenzotriazole (1.1 g, 8.4 mmol) in anhydrous THF (50 mL) was added triethylamine (4.0 mL, 28.0 mmol), and the resulting solution was stirred at rt for 10 min. To the
- 10        resulting reaction solution was then added the appropriate amine or alcohol of the formula X-H (6.7 mmol, 1.2 eq), and this solution was stirred at rt for 24 h. The resulting reaction solution was concentrated via evaporation under reduced pressure, and the residual solid was resuspended in EtOAc (250 mL). This EtOAc suspension was washed with H<sub>2</sub>O (200 mL), NaOH (0.1 N,
- 15        2 x 200 mL), potassium phosphate buffer (50 mM, pH 7, 2 x 200 mL) and H<sub>2</sub>O (200 mL). The organic layer was then dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated via evaporation under reduced pressure until a solid began to precipitate. This mixture was allowed to stand at rt, and the precipitated solid was collected and washed thoroughly with Et<sub>2</sub>O and once with 50% Et<sub>2</sub>O-
- 20        EtOAc to afford the appropriate title compound as a solid. A second crop may be obtained from the mother liquors. If a solid is not obtained by this method, then purification of the extraction residue via column chromatography using silica gel and elution with 5% CH<sub>2</sub>Cl<sub>2</sub>-CH<sub>3</sub>OH afforded the appropriate title compound.


25

Method B: The appropriate 4-amino pyrazolopyridine-5-carboxylic acid (2.8 mmol) was suspended in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (20 mL), and oxalyl chloride (0.72 mL, 8.4 mmol) was added to this mixture followed by 2 drops of DMF. The resulting suspension was stirred at rt for 1 h after the suspension

cleared. The resulting reaction solution was evaporated under reduced pressure, and the residue was redissolved in anhydrous  $\text{CH}_2\text{Cl}_2$  (20 mL). To this solution was added sequentially  $\text{Et}_3\text{N}$  (1.9 mL, 14.0 mmol) and then the appropriate amine or alcohol of the formula X-H (4.2 mmol). The solution  
5 was stirred at rt for 2 hr. The solvent was removed and the resulted residue was suspended in 100 mL EtOAc. The organic was subsequently washed with water, 1 N NaOH, and 1 N HCl. The organic layer was then dried over anhydrous  $\text{Na}_2\text{SO}_4$  and concentrated via evaporation under reduced pressure. The resulted crude products have been purified by either  
10 crystallization (>100 mg scale) or preparative HPLC (<100 mg scale).

Method C: The appropriate 4-amino-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid pentafluorophenyl ester (0.16 mmol) was dissolved in anhydrous THF (2 mL) and triethylamine (49 mg, 0.48 mmol).  
15 The appropriate amine or alcohol of the formula X-H (0.32 mmol, 2.0 eq) was then added, and the reaction solution was stirred at rt for 12 h. The resulting reaction solution was diluted with EtOAc (5 mL), and the organic layer was washed with 0.1 N NaOH (5 mL), potassium phosphate buffer (pH 7) and brine. The organic layer was then dried over anhydrous  $\text{Na}_2\text{SO}_4$  and  
20 concentrated via evaporation under reduced pressure to afford the title compound. Purification of the crude product if necessary can be accomplished by either crystallization or preparative HPLC.


Compounds listed in the following Tables prepared by these  
25 procedures are denoted as methods 1A, 1B, or 1C.

Example 2General Procedure for the Preparation of 4-Aminopyrazolopyridines-5-carboxylates and 4-Aminopyrazolopyridines-5-carboxamides

- 5 To a solution of the appropriate 4-chloro-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid chloride (4.4 mmol) in  $CH_2Cl_2$  was added the appropriate amine or alcohol of the formula X-H (6.6 mmol) and  $Et_3N$  (22.0 mmol) at 0 °C. The reaction was stirred from 0 °C to rt until completion. The solvent was removed via evaporation under reduced pressure, and the residue was
- 10 purified by silica gel chromatography.

- To a solution of the resulting 4-chloropyrazolopyridines-5-carboxamide or ester (0.21 mmol) in ethanol (5 mL) was added an appropriate amine of the formula E-H (0.24 mmol) and  $Et_3N$  (1.4 mmol). The solution was heated at reflux for 2 hr before being diluted with  $CH_2Cl_2$ . The organic layer was
- 15 sequentially washed with 1N HCl, and water, dried, and evaporated under reduced pressure. The residue was purified by silica gel chromatography to afford the title compound.

- Compounds listed in the following Tables prepared by this procedure
- 20 are denoted as method 2.

Example 3General Procedure for the Preparation of 4-Oxypyrazolopyridines-5-carboxylates and 4-Oxypyrazolopyridines-5-carboxamides

25


To a solution of the appropriate 4-chloro-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid chloride (4.4 mmol) in  $\text{CH}_2\text{Cl}_2$  was added the appropriate amine or alcohol of the formula  $\text{X}_1\text{-H}$  (6.6 mmol) and  $\text{Et}_3\text{N}$  (22.0 mmol) at 0 °C. The reaction was stirred from 0 °C to rt until completion. The solvent was removed via evaporation under reduced pressure, and the residue was purified by silica gel chromatography.

To a solution of the resulting 4-chloropyrazolopyridine-5-carboxamide or ester (0.21 mmol) in DMF (2 mL) was added an appropriate alcohol of the formula  $\text{R}_1\text{-OH}$  (0.24 mmol) and  $\text{K}_2\text{CO}_3$  (0.42 mmol). The mixture was stirred at 50 °C for 2 hr before being diluted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was sequentially washed with 1N HCl, and water, dried, and evaporated under reduced pressure. The residue was purified by silica gel chromatography to afford the title compound.

Compounds listed in the following Tables prepared by this procedure are denoted as method 3.

#### Example 4

##### General Procedure for the Preparation of 4-Thiopyrazolopyridines-5-carboxylates and 4-Thiopyrazolopyridines-5-carboxamides




To a solution of the appropriate 4-chloro-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid chloride (4.4 mmol) in  $\text{CH}_2\text{Cl}_2$  was added the appropriate amine or alcohol of the formula  $\text{X}_1\text{-H}$  (6.6 mmol) and  $\text{Et}_3\text{N}$  (22.0 mmol) at 0 °C. The reaction was stirred from 0 °C to rt until completion. The solvent was removed via evaporation under reduced pressure, and the residue was purified by silica gel chromatography.

To a solution of the resulting 4-chloropyrazolopyridine-5-carboxamide or ester (0.21 mmol) in DMF (2 mL) was added an appropriate thiol of the formula  $R_1-SH$  (0.24 mmol) and  $K_2CO_3$  (0.42 mmol). The mixture was stirred at 50 °C for 2 hr before being diluted with  $CH_2Cl_2$ . The organic layer was sequentially washed with 1N HCl, and water, dried, and evaporated under reduced pressure. The residue was purified by silica gel chromatography to afford the title compound.

### Example 5

10 Preparation of 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide




15 Using the procedure of method 1A, triethylamine (4.0 ml, 28.0 mmol) was added to a mixture of 4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid (2.0 g, 5.6 mmol), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (EDAC•HCl, 1.6 g, 8.4 mmol), and 1-hydroxybenzotriazole (1.1 g, 8.4 mmol) in anhydrous THF  
20 (50 ml). The resulting solution was stirred at rt for 10 min. To the resulting reaction solution was then added 4-aminomethylpyridine (0.72 g, 6.7 mmol) and this solution was stirred at rt for 24 h. The resulting reaction solution was

concentrated via evaporation under reduced pressure, and the residual solid was resuspended in EtOAc (250 mL). This EtOAc suspension was washed with H<sub>2</sub>O (200 ml), NaOH (0.1 N, 2 x 200 ml), potassium phosphate buffer (50 mM, pH 7, 2 x 200 ml) and H<sub>2</sub>O (200 mL). The organic layer was then dried  
5 over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated via evaporation under reduced pressure until a solid began to precipitate. This mixture was allowed to stand at rt, and the precipitated solid was collected and washed thoroughly with Et<sub>2</sub>O and once with 50% Et<sub>2</sub>O-EtOAc to afford a white solid (2.2 g, 90%):  
mp: 209-210.5 C; LRMS (m/z) 450; <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ 171.8, 156.4, 153.7,  
10 153.1, 151.7, 150.8, 150.5, 135.7, 132.0, 130.5, 128.3, 124.5, 124.3, 114.2, 105.6, 104.4, 57.4, 49.8, 43.7, 43.6, 15.9. Anal. Calc'd for C<sub>23</sub>H<sub>23</sub>C1N<sub>6</sub>O<sub>2</sub>: C, 61.26; H, 5.14; N, 18.64; Cl, 7.86. Found: C, 61.03; H, 5.22; N, 18.66; Cl, 8.02.

Table

The following compounds of formula I were prepared wherein R<sub>3</sub> is

















ethyl, R<sub>4</sub> is hydrogen, Y is nitrogen, Z is CH, and E<sub>1</sub> is








5








Ex	Name	X <sub>1</sub>	PURITY (%)	HPLC (retention time, minutes)	OTHER DATA
6	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[1-(phenylmethyl)-4-piperidinylmethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	100	3.47	m/z (M+H) 489
7	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-(3-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	96	3.28	m/z (M+H) 407
8	(R)-4-[(1-Cyclohexylethyl)amino]-N-(2,3-dihydro-1H-indan-2-yl)-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	91	4.53	m/z (M+H) 432
9	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[(4-hydroxy-3-methoxyphenyl)methyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	91	3.91	m/z (M+H) 452

















10	(1R)-4-[(1-Cyclohexylethyl)amino]-N-[(2,3-dihydro-1,4-benzodioxin-2-yl)methyl]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	93	4.56	m/z (M+H) 464
11	(R)-N-(1H-Benzimidazol-2-ylmethyl)-4-[(1-cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	90	3.58	m/z (M+H) 446
12	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[3-(2-oxo-1-pyrrolidinyl)propyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	90	3.72	m/z (M+H) 441
13	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[(5-methyl-2-furanyl)methyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	94	4.28	m/z (M+H) 410
14	(R)-N-[(2-Chlorophenyl)methyl]-4-[(1-cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	93	4.52	m/z (M+H) 440
15	(1R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[3-(2-methyl-1-piperidinyl)propyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	88	3.25	m/z (M+H) 455
16	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[2-(2-pyridinyl)ethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	88	3.19	m/z (M+H) 421








17	(1R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-(hexahydro-2-oxo-1H-azepin-3-yl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	84	3.84	m/z (M+H) 427
18	[R-(R*,R*)]-a-[[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]amino]benzeneacetic acid ethyl ester	 (METHOD 1C)	93	4.43	m/z (M+H) 478
19	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[2-(3-pyridinyl)ethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	93	3.20	m/z (M+H) 420
20	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[[4-(1,2,3-thiadiazol-4-yl)phenyl]methyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	85	4.26	m/z (M+H) 490
21	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-(2-thienylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	93	4.27	m/z (M+H) 412
22	(1R)-N-(1-Azabicyclo[2.2.2]octan-3-yl)-4-[(1-cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	85	3.21	m/z (M+H) 425
23	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-(3-1H-imidazol-1-ylpropyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	94	3.11	m/z (M+H) 424








24	(R)-4-[[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-yl]carbonyl]-amino]-1-piperidinecarboxylic acid ethyl ester	 (METHOD 1C)	90	4.23	m/z (M+H) 471
25	[1S-[1a(S*),2b)]-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-(2-phenylcyclopropyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	95	4.54	m/z (M+H) 432
26	(R)-4-[(1-Cyclohexylethyl)amino]-N-[(2,6-difluorophenyl)methyl]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	93	4.3	m/z (M+H) 441
27	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-(phenylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	95	4.33	m/z (M+H) 406
28	(R)-4-[(1-Cyclohexylethyl)amino]-N-[6-(dimethylamino)hexyl]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	84	3.29	m/z (M+H) 443
29	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[2-(2-methoxyphenyl)ethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	91	4.45	m/z (M+H) 449
30	[R-(R*,R*)]-N-(1-Cyclohexylethyl)-4-[(1-cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	92	4.66	m/z (M+H) 426

31	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[2-(4-methoxyphenyl)ethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	95	4.38	m/z (M+H) 449
32	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-(4-hydroxybutyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	94	3.67	m/z (M+H) 388
33	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[2-(5-methoxy-1H-indol-3-yl)ethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	88	4.18	m/z (M+H) 488
34	[1S-[1a,2a(S*),4a]]-N-(Bicyclo[2.2.1]heptan-2-yl)-4-[(1-cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	88	4.47	m/z (M+H) 410
35	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-(2-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	98	3.42	m/z (M+H) 407
36	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[(1-ethyl-2-pyrrolidinyl)methyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	84	3.28	m/z (M+H) 427
37	(R)-N-[2-(Acetylamino)ethyl]-4-[(1-cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	93	3.52	m/z (M+H) 401







38	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	89	3.25	m/z (M+H) 407
39	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[2-(2-thienyl)ethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	92	4.36	m/z (M+H) 425
40	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[2-(1-methyl-2-pyrrolidinyl)ethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	94	3.19	m/z (M+H) 427
41	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[3-(4-morpholinyl)propyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	85	3.09	m/z (M+H) 442
42	(R)-4-[(1-Cyclohexylethyl)amino]-N-[(2,4-dimethoxyphenyl)methyl]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	93	4.3	m/z (M+H) 466
43	(R)-4-[(1-Cyclohexylethyl)amino]-N-[2-(3,4-dimethoxyphenyl)ethyl]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	94	4.19	m/z (M+H) 480
44	(R)-4-[(1-Cyclohexylethyl)amino]-N-[(3,4-difluorophenyl)methyl]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	93	4.50	m/z (M+H) 442








45	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[2-(1-pyrrolidinyl)ethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	91	3.15	m/z (M+H) 413
46	(R)-N-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-yl]carbonyl]glycine ethyl ester	 (METHOD 1C)	90	4.01	m/z (M+H) 401
47	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[1-(phenylmethyl)-3-pyrrolidinyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	92	3.54	m/z (M+H) 475
48	(R)-N-[2-[4-(Aminosulfonyl)phenyl]ethyl]-4-[(1-cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	93	3.76	m/z (M+H) 498
49	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[(3,4,5-trimethoxyphenyl)methyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	93	4.17	m/z (M+H) 496
50	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-(6-hydroxyhexyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	94	3.96	m/z (M+H) 416
51	(1R)-4-[(1-Cyclohexylethyl)amino]-N-(2,3-dihydroxypropyl)-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	95	3.39	m/z (M+H) 390








52	[S-(R*,S*)]-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-2-pyrrolidinecarboxamide	 (METHOD 1C)	94	3.18	m/z (M+H) 413
53	(R)-1-(1,3-Benzodioxol-5-ylmethyl)-4-[[4-[(1-cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]piperazine	 (METHOD 1C)	93	3.27	m/z (M+H) 519
54	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-(2-hydroxyethyl)-N-propyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	96	3.53	m/z (M+H) 402
55	(R)-4-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-1-piperazinecarboxylic acid ethyl ester	 (METHOD 1C)	95	3.90	m/z (M+H) 457
56	(R)-4-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-morpholine	 (METHOD 1C)	100	3.51	m/z (M+H) 386
57	(R)-8-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-oxo-1-phenyl-1,3,8-triazaspiro[4.5]decane	 (METHOD 1C)	95	4.18	m/z (M+H) 530
58	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-methyl-N-(1-methyl-4-piperidinyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	92	2.79	m/z (M+H) 427









59	[S-(R*,S*)]-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-2-(hydroxymethyl)pyrrolidine	 (METHOD 1C)	85	3.48	m/z (M+H) 400
60	(R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-(phenylmethyl)piperidine	 (METHOD 1C)	92	4.62	m/z (M+H) 474
61	[S-(R*,S*)]-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-2-[[2,6-dimethylphenyl]amino]methylpyrrolidine	 (METHOD 1C)	93	4.05	m/z (M+H) 503
62	(R)-4-[(1-Cyclohexylethyl)amino]-N-[2-(diethylamino)ethyl]-1-ethyl-N-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	97	2.90	m/z (M+H) 429
63	(R)-4-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]thiomorpholine	 (METHOD 1C)	90	3.87	m/z (M+H) 402
64	(R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]piperidine	 (METHOD 1C)	95	3.92	m/z (M+H) 384
65	(1R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-3-piperidinecarboxylic acid ethyl ester	 (METHOD 1C)	94	3.99	m/z (M+H) 456








66	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-propyl-N-[2-(2-pyridinyl)ethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	100	3.48	m/z (M+H) 463
67	(1R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-3-(hydroxymethyl)piperidine	 (METHOD 1C)	100	3.50	m/z (M+H) 414
68	(R)-2-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-1,2,3,4-tetrahydro-6,7-dimethoxyisoquinoline	 (METHOD 1C)	96	3.96	m/z (M+H) 492
69	(R)-4-(4-Chlorophenyl)-1-[[4-[(1-cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-1,2,3,6-tetrahydropyridine	 (METHOD 1C)	95	4.69	m/z (M+H) 492
70	(1R)-4-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-1-(4-methoxyphenyl)-2-methylpiperazine	 (METHOD 1C)	96	3.91	m/z (M+H) 505
71	(R)-1-[Bis(4-fluorophenyl)methyl]-4-[[4-[(1-cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl-piperazine	 (METHOD 1C)	93	4.53	m/z (M+H) 587


72	(R)-4-[(1-Cyclohexylethyl)amino]-N-[2-(dimethylamino)ethyl]-1-ethyl-N-(phenylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	76	3.61	m/z (M+H) 477
73	(R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-phenylpiperazine	 (METHOD 1C)	93	4.31	m/z (M+H) 461
74	(R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-(2-hydroxyethyl)piperazine	 (METHOD 1C)	92	2.89	m/z (M+H) 429
75	(R)-1-(2-Chlorophenyl)-4-[[4-[(1-cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]piperazine	 (METHOD 1C)	94	4.63	m/z (M+H) 495
76	(1R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-3-piperidinecarboxamide	 (METHOD 1C)	94	3.31	m/z (M+H) 427
77	(R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-hydroxypiperidine	 (METHOD 1C)	98	3.34	m/z (M+H) 400
78	(R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-methylpiperidine	 (METHOD 1C)	94	4.17	m/z (M+H) 398


79	(R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-(3,4-dichlorophenyl)piperazine	 (METHOD 1C)	92	4.75	m/z (M+H) 529
80	[S-(R*,S*)]-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-2-(methoxymethyl)pyrrolidine	 (METHOD 1C)	80	3.85	m/z (M+H) 414
81	(R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-(3-methoxyphenyl)piperazine	 (METHOD 1C)	98	4.32	m/z (M+H) 491
82	(R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-(4-methoxyphenyl)piperazine	 (METHOD 1C)	95	4.14	m/z (M+H) 491
83	(1R)-N-[1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-3-pyrrolidinyl]acetamide	 (METHOD 1C)	95	3.31	m/z (M+H) 427
84	(R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]hexahydro-4-methyl-1H-1,4-diazepine	 (METHOD 1C)	94	2.76	m/z (M+H) 413
85	(1R)-4-[(1-Cyclohexylethyl)amino]-N,1-diethyl-N-(1-ethyl-3-pyrrolidinyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	90	2.93	m/z (M+H) 441






86	(R,Z)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-(3-phenyl-2-propenyl)piperazine	 (METHOD 1C)	96	3.52	m/z (M+H) 501
87	(1R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-hydroxy-4-(phenylmethyl)piperidine	 (METHOD 1C)	93	4.18	m/z (M+H) 490
88	(R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-piperidinecarboxamide	 (METHOD 1C)	93	3.22	m/z (M+H) 427
89	(R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-(2-pyridinyl)piperazine	 (METHOD 1C)	94	3.01	m/z (M+H) 462
90	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-methyl-N-[2-(2-pyridinyl)ethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	94	2.92	m/z (M+H) 435
91	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N,N-bis(2-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	92	3.43	m/z (M+H) 498
92	(R)-1-Acetyl-4-[[4-[(1-cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]piperazine	 (METHOD 1C)	97	3.36	m/z (M+H) 427
93	(R)-1-[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-(2-fluorophenyl)piperazine	 (METHOD 1C)	92	4.41	m/z (M+H) 479

94	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-methyl-N-[2-(4-pyridinyl)ethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	92	2.91	m/z (M+H) 435
95	(1R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-methyl-N-[1-(phenylmethyl)-3-pyrrolidinyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1C)	93	3.36	m/z (M+H) 489
96	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-3-pyridinyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	96	3.57	m/z (M+H) 393
97	(R)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[2-(4-morpholinyl)ethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	98	3.23	m/z (M+H) 429
98	4-[[[(R)-1-Cyclohexylethyl]amino]-1-ethyl-N-[1-(4-hydroxyphenyl)ethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	99	3.78 & 3.87	m/z (M+H) 436







The following compounds of formula I were prepared wherein R<sub>3</sub> is




Ex.	Name	X <sub>2</sub>	PURITY (%)	HPLC (retention time, minutes)	OTHER DATA
99	(S)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	100	3.27	m/z (M+H) 407

100	(S)-4-[[[4-[(1-Cyclohexylethyl)amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]amino]-1-piperidinecarboxylic acid ethyl ester	 (METHOD 1A)	89	4.07	m/z (M+H) 471
101	(S)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-(3-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	100	3.28	m/z (M+H) 407
102	(1S)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-(hexahydro-2-oxo-1H-azepin-3-yl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	97	3.71	m/z (M+H) 427
103	(S)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[3-(2-oxo-1-pyrrolidinyl)propyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	90	3.65	m/z (M+H) 441
104	(S)-4-[(1-Cyclohexylethyl)amino]-1-ethyl-N-[(4-hydroxy-3-methoxyphenyl)methyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	97	3.81	m/z (M+H) 452

The following compounds of formula I were prepared wherein R<sub>3</sub> is ethyl, R<sub>4</sub> is hydrogen, Y is nitrogen, and Z is CH:






Ex.	Name	E <sub>1</sub>	X <sub>1</sub>	PURITY (%)	HPLC (retention time, minutes)	OTHER DATA
105	4-[[[4-Morpholinyl)-ethyl]amino]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide		 Method 2	99	0.80	m/z (M+H) 410
106	4-[(N-(3-1H-Imidazol-1-yl)propyl)amino]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide		 Method 2	98	1.20	m/z (M+H) 405
107	4-[[[3-Chloro-4-methoxyphenyl)methyl]hydroxyl]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide		 Method 2	99	2.81	m/z (M+H) 452







The following compounds of formula II were prepared wherein R<sub>3</sub> is ethyl, R<sub>4</sub> is hydrogen, Y is nitrogen, Z is CH, and E<sub>2</sub> is














Ex.	Name	X <sub>2</sub>	PURITY (%)	HPLC (retention time, minutes)	OTHER DATA
108	4-[[[4-[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]amino]-1-piperidine-carboxylic acid ethyl ester	 (METHOD 1B)	96	3.62	mp 184-185 °C; Anal. Calcd for C <sub>25</sub> H <sub>31</sub> ClN <sub>6</sub> O <sub>4</sub> : C, 58.30; H, 6.07; N, 16.32; Cl, 6.88. Found: C, 58.14; H, 6.18; N, 16.22; Cl, 6.58.
109	4-[[[3-(2-oxo-1-pyrrolidinyl)propyl]amino]-1-ethyl-N-[3-(2-oxo-1-pyrrolidinyl)propyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	98	3.2	mp 149-150 °C; Anal. Calcd for C <sub>24</sub> H <sub>29</sub> ClN <sub>6</sub> O <sub>3</sub> : C, 59.44; H, 6.03; N, 17.33; Cl, 7.31. Found: C, 59.54; H, 5.89; N, 17.38; Cl, 7.37.













110	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(2-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	98	2.95	mp 176 - 177 °C; Anal. Calcd for C <sub>23</sub> H <sub>23</sub> ClN <sub>6</sub> O <sub>2</sub> : C, 61.26; H, 5.14; N, 18.64; Cl, 7.86. Found: C, 60.99; H, 5.00; N, 18.71; Cl, 7.92.
111	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(3-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	98	2.90	mp 198 °C NMR C <sup>13</sup> H <sup>1</sup> m/z (M+H) 451
112	4-[[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]amino]-1-piperidinecarboxylic acid ethyl ester	 (METHOD 1A)	98	3.70	mp 182 - 184 °C NMR C <sup>13</sup> H <sup>1</sup> m/z (M+H) 515
113	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(hexahydro-2-oxo-1H-azepin-3-yl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	93	3.30	mp 214 - 216 °C NMR C <sup>13</sup> H <sup>1</sup> m/z (M+H) 471
114	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[(4-hydroxy-3-methoxyphenyl)methyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	99	3.45	mp 119 - 121 °C NMR C <sup>13</sup> H <sup>1</sup> m/z (M+H) 496







115	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[[4-(1,2,3-thiadiazol-4-yl)phenyl]methyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	97	3.30	mp 194 - 196 °C NMR C <sup>13</sup> H <sup>1</sup> m/z (M+H) 534
116	N-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]glycine ethyl ester	 (METHOD 1A)	95	3.40	mp 149 - 152 °C NMR C <sup>13</sup> H <sup>1</sup> m/z (M+H) 446
117	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(3-1H-imidazol-1-ylpropyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	99	2.90	mp 108 - 110 °C NMR C <sup>13</sup> H <sup>1</sup> m/z (M+H) 468
118	N-[(3-Chloro-4-methoxyphenyl)methyl]-4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	93	3.80	NMR C <sup>13</sup> H <sup>1</sup> m/z (M+H) 514
119	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N,N-bis(2-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	92	3.00	NMR C <sup>13</sup> H <sup>1</sup> m/z (M+H) 542
120	(R)-α-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]amino]benzeneacetic acid ethyl ester	 (METHOD 1A)	94	3.80	NMR C <sup>13</sup> H <sup>1</sup> m/z (M+H) 522





121	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[1-(phenylmethyl)-4-piperidiny]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	96	3.10	NMR C <sup>13</sup> H <sup>1</sup> m/z (M+H) 533
122	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[3-(4-morpholinyl)propyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	98	2.80	NMR C <sup>13</sup> H <sup>1</sup> m/z (M+H) 487
123	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-N-[2-(dimethylamino)ethyl]-1-ethyl-N-(phenylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	88	3.10	NMR C <sup>13</sup> H <sup>1</sup> m/z (M+H) 521
124	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[(4-pyridinyl-1-oxide)methyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	97	3.20	NMR C <sup>13</sup> H <sup>1</sup> m/z (M+H) 467
125	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(2-furanylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	100	3.73	m/z (M+H) 440
126	N-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-L-valine ethyl ester	 (METHOD 1B)	92	4.06	m/z (M+H) 488

127	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-N-(cyanomethyl)-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	92	3.40	m/z (M+H) 399
128	N2-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-N6-[(phenylmethoxy)carbonyl]-L-lysine methyl ester	 (METHOD 1B)	82	4.07	m/z (M+H) 637
129	N-[2-(Acetylamino)ethyl]-4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	100	3.12	m/z (M+H) 445
130	N-[N-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-L-alanyl]glycine methyl ester	 (METHOD 1B)	100	3.17	m/z (M+H) 503
131	[4-[[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]amino]butyl]carbamic acid 1,1-dimethylethyl ester	 (METHOD 1B)	96	2.79	m/z (M+H) 531



132	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(1,2,3,4-tetrahydro-1-naphthalenyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	98	4.33	m/z (M+H) 490
133	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(tetrahydro-2-furanylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	92	3.53	m/z (M+H) 444
134	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-N-[[4-(dimethylamino)phenyl]methyl]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	100	3.09	m/z (M+H) 493
135	N-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-L-alanine methyl ester	 (METHOD 1B)	97	3.53	m/z (M+H) 446
136	(S)-2-(Acetylamino)-6-[[[4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]amino]hexanamide	 (METHOD 1B)	100	3.25	m/z (M+H) 544







137	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[[4-fluoro-2-(trifluoromethyl)phenyl]methyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	96	4.39	m/z (M+H) 536
138	(4S-cis)-4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(2,2-dimethyl-4-phenyl-1,3-dioxan-5-yl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	88	4.07	m/z (M+H) 550
139	N-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-L-methionine methyl ester	 (METHOD 1B)	85	3.89	m/z (M+H) 506
140	2-[[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]amino]propanedioic acid diethyl ester	 (METHOD 1B)	74	3.95	m/z (M+H) 518
141	4-[[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]amino]butanoic acid methyl ester	 (METHOD 1B)	81	3.56	m/z (M+H) 460






142	N-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-L-alanine ethyl ester	 <p>(METHOD 1B)</p>	83	3.73	m/z (M+H) 460
143	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(2-oxo-2-phenylethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 <p>(METHOD 1B)</p>	100	3.83	m/z (M+H) 478
144	1-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-(2-furanylcarbonyl)piperazine	 <p>(METHOD 1B)</p>	87	3.17	m/z (M+H) 523
145	(S)-4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(tetrahydro-2-oxo-3-furanyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamid	 <p>(METHOD 1B)</p>	96	3.24	m/z (M+H) 444
146	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-4-morpholinyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 <p>(METHOD 1B)</p>	100	3.27	m/z (M+H) 445
147	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[2-(4-morpholinyl)ethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 <p>(METHOD 1B)</p>	100	2.77	m/z (M+H) 473

148	N-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-L-asparagine 1,1-dimethylethyl ester	 (METHOD 1B)	77	3.69	m/z (M+H) 531
149	(S)-4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[2-oxo-3-[2-oxo-2-(1-pyrrolidinyl)ethyl]cycloheptyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	98	3.52	m/z (M+H) 582
150	(S)-α-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]amino]-4-hydroxybenzeneacetic acid	 (METHOD 1A)	72	3.15	m/z (M+H) 510
151	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(4-piperidinyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	92	2.82	m/z (M+H) 443









152	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxylate	 METHOD 2 (also made using the procedure of Method 1A)	95	3.15	m/z (M+H) 452
153	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[(3-(4-morpholinyl)ethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxylate	 METHOD 2	95	2.97	m/z (M+H) 474




154	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(4-pyridinylethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (Method 1A)	97	2.89	m/z (M+H) 465
155	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(2-pyridinylethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (Method 1A)	90	2.90	m/z (M+H) 465
156	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[3-(1-N-methylpiperizinyl)propyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (Method 1A)	98	2.60	m/z (M+H) 500
157	(S)-4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(2-tetrahydrofurylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (Method 1A)	99	3.70	m/z (M+H) 444
158	(R)-4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(2-tetrahydrofurylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (Method 1A)	94	3.6	m/z (M+H) 444
159	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[4-(2-chloropyridinyl)methyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (Method 1A)	90	3.96	m/z (M+H) 485

160	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[4-(2,6-dichloropyridinyl)methyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (Method 1A)	97	3.90	m/z (M+H) 519, 521
161	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(5-tetrazoyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (Method 1A)	83	3.10	m/z (M+H)
162	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[4-(morpholinylethyl)]-1H-pyrazolo[3,4-b]pyridine-5-carboxylate	 (Method 1A)	85	3.00	m/z (M+H) 474
163	4-[N-(methylpropylate)amino]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (Method 3)	88	1.90	m/z (M+H) 397
164	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(2-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxylate	 (Method 2)	95	3.40	m/z (M+H) 452


The following compounds of formula II were prepared wherein R<sub>3</sub> is ethyl, R<sub>4</sub> is hydrogen, Y is nitrogen, and Z is CH<sub>2</sub>:



5






Ex.	Name	E <sub>2</sub>	X <sub>2</sub>	PURITY (%)	HPLC (retention time, minutes)	OTHER DATA
165	4-[[[(3-Fluoro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide		 Method 2	97	2.65	m/z (M+H) 435
166	4-[[[(3,5-Dichloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide		 Method 2	98	3.18	m/z (M+H) 486
167	4-[[[(4-Fluorophenyl)methyl]amino]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide		 Method 2	99	2.67	m/z (M+H) 405







168	4-[[[4-Methoxy)butyl]amino]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	$\text{—N—}(\text{CH}_2)_4\text{O—Me}$ H	 Method 2	98	2.20	m/z (M+H) 383
169	4-[[[(N, N'-3-Dimethylamino)propyl]amino]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	$\text{—N—}(\text{CH}_2)_3\text{N(Me)}_2$ H	 Method 2	95	0.93	m/z (M+H) 382
170	4-[(1,3-Benzodioxol-5-ylmethyl)amino]-1-ethyl-N-[3-(2-oxo-1-pyrrolidinyl)propyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 H	$\text{—N—}(\text{CH}_2)_3\text{N}$ H	91	3.07	m/z (M+H) 465

The following compounds of formula III were prepared wherein R<sub>3</sub> is ethyl, R<sub>4</sub> is hydrogen, Y is nitrogen, Z is CH<sub>2</sub>, and E<sub>1</sub> is







Ex.	Name	X <sub>3</sub>	PURITY (%)	HPLC (retention time, minutes)	OTHER DATA
171	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(6-hydroxyhexyl)-1H-pyrazolo[3,4-b]pyridine-5-	$-\text{NH}-(\text{CH}_2)_6-\text{OH}$ (METHOD 1A)	97	3.40	mp 117 - 118 °C; NMR C <sup>13</sup> H <sup>1</sup> m/z (M+H) 460.
172	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-N-[4-(diethylamino)-1-methylbutyl]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	95	3.03	m/z (M+H) 501
173	N-[2-[Bis(1-methylethyl)amino]ethyl]-4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	85	3.06	m/z (M+H) 487

174	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-N-[2,2-dimethyl-3-(dimethylamino)propyl]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 <p>(METHOD 1B)</p>	95	2.85	m/z (M+H) 473
175	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[1-(methoxymethyl)propyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 <p>(METHOD 1B)</p>	100	3.67	m/z (M+H) 446
176	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[3-(1-methylethoxy)propyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 <p>(METHOD 1B)</p>	83	3.9	m/z (M+H) 460
177	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(4-methoxybutyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 <p>(METHOD 1B)</p>	100	3.64	m/z (M+H) 446
178	(R)-1-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-3-(dimethylamino)pyrrolidine	 <p>(METHOD 1B)</p>	99	2.41	m/z (M+H) 457


179	(S)-1-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-3-(dimethylamino)pyrrolidine	 (METHOD 1B)	88	2.40	m/z (M+H) 457
180	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-methoxy-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	89	3.38	m/z (M+H) 390
181	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-N-[2-(dimethylamino)ethyl]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	96	2.78	m/z (M+H) 431
182	1-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-4-methylpiperazine	 (METHOD 1B)	97	2.53	m/z (M+H) 443
183	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(3-hydroxypropyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	94	3.01	m/z (M+H) 418
184	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(4-hydroxybutyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	94	3.11	m/z (M+H) 432



185	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(5-hydroxypentyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	88	3.22	m/z (M+H) 446
186	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(2-hydroxyethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1B)	97	2.9	m/z (M+H) 404
187	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(4-hydroxypiperidinyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	97	2.80	m/z (M+H) 444
188	4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[3-(1,2-dihydroxypropyl)]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide	 (METHOD 1A)	95	2.97	m/z (M+H) 434


What is claimed is:

1. A compound of the formulas  
(I)



5

(II)



- including a pharmaceutically acceptable salt thereof  
10 wherein:

E<sub>1</sub> is -O-R<sub>1</sub>, -S-R<sub>1</sub>, -NH-A<sub>1</sub>-cycloalkyl, -NH-A<sub>1</sub>-substituted cycloalkyl, -NH-A<sub>1</sub>-heterocyclo, or -NH-A<sub>1</sub>-heteroaryl;

E<sub>2</sub> is -NH-A<sub>1</sub>-alkoxy, -NH-A<sub>1</sub>-CO<sub>2</sub>alkyl,

- 15  $\text{—NH—A}_1\text{—N} \begin{matrix} \text{R}_{15} \\ \text{R}_{16} \end{matrix}$ , -NH-A<sub>1</sub>-aryl, or -NH-A<sub>1</sub>-substituted aryl;

R<sub>1</sub> is -A<sub>1</sub>-cycloalkyl, -A<sub>1</sub>-substituted cycloalkyl, -A<sub>1</sub>-alkoxy,  $\text{—A}_1\text{—N} \begin{matrix} \text{R}_{15} \\ \text{R}_{16} \end{matrix}$ , -A<sub>1</sub>-aryl, -A<sub>1</sub>-substituted aryl,  
20 -A<sub>1</sub>-heterocyclo, or -A<sub>1</sub>-heteroaryl;

$X_1$  is  $-O-A_1-R_2$ ,  $-O-R_9$ ,  $-N(R_9)(R_{10})$ ,  $\begin{array}{c} -N-A_2-R_2 \\ | \\ R_5 \end{array}$ , a

monocyclic ring  $\begin{array}{c} -N \\ \diagup \quad \diagdown \\ (R_{21})_n \quad (R_{22})_m \end{array}$ , a fused bicyclic ring

$\begin{array}{c} -N \quad B_1 \\ \diagup \quad \diagdown \\ (R_{21})_n \quad (R_{22})_m \end{array}$ , or a spiro ring  $\begin{array}{c} -N \quad C \quad B_2 \\ \diagup \quad \diagdown \\ (R_{21})_n \quad (R_{22})_m \end{array}$ ;

$X_2$  is  $-O-A_1-R_{25}$ ,  $\begin{array}{c} -N-A_2-R_{25} \\ | \\ R_5 \end{array}$ , a monocyclic ring

5  $\begin{array}{c} -N \\ \diagup \quad \diagdown \\ (R_{21})_n \quad R_{22} \end{array}$ , a fused bicyclic ring  $\begin{array}{c} -N \quad B_1 \\ \diagup \quad \diagdown \\ (R_{21})_n \quad (R_{22})_m \end{array}$ ,

or a spiro ring  $\begin{array}{c} -N \quad C \quad B_2 \\ \diagup \quad \diagdown \\ (R_{21})_n \quad (R_{22})_m \end{array}$ ;

$X_3$  is  $-O-R_9$ ,  $-O-A_1-O-R_9$ ,  $-N(R_9)(R_{10})$ ,  $\begin{array}{c} -N-A_2-O-R_9 \\ | \\ R_5 \end{array}$ ,

$\begin{array}{c} -N-A_1-N(R_9)(R_{10}) \\ | \\ R_5 \end{array}$ , or a monocyclic ring  $\begin{array}{c} -N \\ \diagup \quad \diagdown \\ (R_{21})_n \end{array}$ ;

10

$A_1$  is an alkylene or substituted alkylene bridge of 1 to 10 carbons;

$Y$  is nitrogen or  $C(R_6)$ ;

15  $Z$  is nitrogen or  $C(R_7)$  with the proviso that at least one of  $Y$  or  $Z$  is nitrogen;

$R_3$  is hydrogen, alkyl, cycloalkyl, substituted cycloalkyl, substituted alkyl,  $-A_1$ -aryl,  $-A_1$ -substituted aryl,  $-A_1$ -cycloalkyl, or  $-A_1$ -substituted cycloalkyl;

R<sub>6</sub> and R<sub>7</sub> are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, -A<sub>1</sub>-cycloalkyl, -A<sub>1</sub>-substituted cycloalkyl, -A<sub>1</sub>-aryl, A<sub>1</sub>-substituted aryl, 5 -A<sub>1</sub>-heterocyclo, and A<sub>1</sub>-heteroaryl;

R<sub>4</sub> is hydrogen, -N(R<sub>12</sub>)(R<sub>13</sub>), -OR<sub>12</sub> or 1- or 3-imidazolyl;

A<sub>2</sub> is a direct bond, an alkylene or substituted alkylene bridge of 1 to 10 carbons, an alkenyl or 10 substituted alkenyl bridge of 2 to 10 carbons having one or more double bonds, or an alkynyl or substituted alkynyl bridge of 2 to 10 carbons having one or more triple bonds;


R<sub>2</sub> is cycloalkyl, substituted cycloalkyl, aryl, 15 substituted aryl, heterocyclo, heteroaryl, cycloalkyl-A<sub>3</sub>-cycloalkyl, cycloalkyl-A<sub>3</sub>-substituted cycloalkyl, cycloalkyl-A<sub>3</sub>-aryl, cycloalkyl-A<sub>3</sub>-substituted aryl, cycloalkyl-A<sub>3</sub>-heterocyclo, cycloalkyl-A<sub>3</sub>-heteroaryl, substituted cycloalkyl-A<sub>3</sub>-cycloalkyl, substituted 20 cycloalkyl-A<sub>3</sub>-substituted cycloalkyl, substituted cycloalkyl-A<sub>3</sub>-aryl, substituted cycloalkyl-A<sub>3</sub>-substituted aryl, substituted cycloalkyl-A<sub>3</sub>-heterocyclo, substituted cycloalkyl-A<sub>3</sub>-heteroaryl, aryl-A<sub>3</sub>-cycloalkyl, aryl-A<sub>3</sub>-substituted cycloalkyl, aryl-A<sub>3</sub>-aryl, aryl-A<sub>3</sub>-substituted 25 aryl, aryl-A<sub>3</sub>-heterocyclo, aryl-A<sub>3</sub>-heteroaryl, substituted aryl-A<sub>3</sub>-cycloalkyl, substituted aryl-A<sub>3</sub>-substituted cycloalkyl, substituted aryl-A<sub>3</sub>-aryl, substituted aryl-A<sub>3</sub>-substituted aryl, substituted aryl-A<sub>3</sub>-heterocyclo, substituted aryl-A<sub>3</sub>-heteroaryl, heterocyclo-A<sub>3</sub>-cycloalkyl, 30 heterocyclo-A<sub>3</sub>-substituted cycloalkyl, heterocyclo-A<sub>3</sub>-aryl, heterocyclo-A<sub>3</sub>-substituted aryl, heterocyclo-A<sub>3</sub>-heterocyclo, heterocyclo-A<sub>3</sub>-heteroaryl, heteroaryl-A<sub>3</sub>-cycloalkyl, heteroaryl-A<sub>3</sub>-substituted cycloalkyl, heteroaryl-A<sub>3</sub>-aryl, heteroaryl-A<sub>3</sub>-heterocyclo, heteroaryl-A<sub>3</sub>-heteroaryl, cyano, -OR<sub>9</sub>, -SR<sub>9</sub>, -(C=O)R<sub>9</sub>, -N(R<sub>9</sub>)(R<sub>10</sub>), 35

-CO<sub>2</sub>R<sub>9</sub>, -(C=O)N(R<sub>12</sub>)(R<sub>13</sub>), -SO<sub>2</sub>N(R<sub>12</sub>)(R<sub>13</sub>), -NR<sub>11</sub>(C=O)R<sub>19</sub>,  
 -NR<sub>11</sub>(C=O)N(R<sub>12</sub>)(R<sub>13</sub>), -O-(C=O)N(R<sub>12</sub>)(R<sub>13</sub>) provided that A<sub>2</sub>  
 is not a direct bond, -NR<sub>11</sub>CO<sub>2</sub>R<sub>19</sub>, -(C=O)N(R<sub>11</sub>)CH<sub>2</sub>CO<sub>2</sub>R<sub>19</sub>,  
 nitrogen when A<sub>2</sub> is alkynyl ending in a triple bond, or  
 5 NH when A<sub>2</sub> is alkenyl ending in a double bond;

R<sub>25</sub> is cycloalkyl, substituted cycloalkyl, aryl,  
 substituted aryl, heterocyclo, heteroaryl, cycloalkyl-A<sub>3</sub>-  
 cycloalkyl, cycloalkyl-A<sub>3</sub>-substituted cycloalkyl,  
 cycloalkyl-A<sub>3</sub>-aryl, cycloalkyl-A<sub>3</sub>-substituted aryl,  
 10 cycloalkyl-A<sub>3</sub>-heterocyclo, cycloalkyl-A<sub>3</sub>-heteroaryl,  
 substituted cycloalkyl-A<sub>3</sub>-cycloalkyl, substituted  
 cycloalkyl-A<sub>3</sub>-substituted cycloalkyl, substituted  
 cycloalkyl-A<sub>3</sub>-aryl, substituted cycloalkyl-A<sub>3</sub>-substituted  
 aryl, substituted cycloalkyl-A<sub>3</sub>-heterocyclo, substituted  
 15 cycloalkyl-A<sub>3</sub>-heteroaryl, aryl-A<sub>3</sub>-cycloalkyl, aryl-A<sub>3</sub>-  
 substituted cycloalkyl, aryl-A<sub>3</sub>-aryl, aryl-A<sub>3</sub>-substituted  
 aryl, aryl-A<sub>3</sub>-heterocyclo, aryl-A<sub>3</sub>-heteroaryl, substituted  
 aryl-A<sub>3</sub>-cycloalkyl, substituted aryl-A<sub>3</sub>-substituted  
 cycloalkyl, substituted aryl-A<sub>3</sub>-aryl,  
 20 substituted aryl-A<sub>3</sub>-substituted aryl, substituted aryl-A<sub>3</sub>-  
 heterocyclo, substituted aryl-A<sub>3</sub>-heteroaryl, heterocyclo-  
 A<sub>3</sub>-cycloalkyl, heterocyclo-A<sub>3</sub>-substituted cycloalkyl,  
 heterocyclo-A<sub>3</sub>-aryl, heterocyclo-A<sub>3</sub>-substituted aryl,  
 heterocyclo-A<sub>3</sub>-heterocyclo, heterocyclo-A<sub>3</sub>-heteroaryl,  
 25 heteroaryl-A<sub>3</sub>-cycloalkyl, heteroaryl-A<sub>3</sub>-substituted  
 cycloalkyl, heteroaryl-A<sub>3</sub>-aryl, heteroaryl-A<sub>3</sub>-substituted  
 aryl, heteroaryl-A<sub>3</sub>-heterocyclo, heteroaryl-A<sub>3</sub>-heteroaryl,  
 cyano, -S-R<sub>9</sub>, -(C=O)R<sub>11</sub>, -CO<sub>2</sub>R<sub>19</sub>, -(C=O)N(R<sub>12</sub>)(R<sub>13</sub>),  
 -SO<sub>2</sub>N(R<sub>12</sub>)(R<sub>13</sub>), -NR<sub>9</sub>(C=O)R<sub>10</sub>, -NR<sub>11</sub>(C=O)N(R<sub>12</sub>)(R<sub>13</sub>),  
 30 -O-(C=O)N(R<sub>12</sub>)(R<sub>13</sub>) provided that A<sub>2</sub> is not a direct bond,  
 -NR<sub>11</sub>CO<sub>2</sub>R<sub>19</sub>, -(C=O)N(R<sub>11</sub>)CH<sub>2</sub>CO<sub>2</sub>R<sub>19</sub>, nitrogen when A<sub>2</sub> is  
 alkynyl ending in a triple bond, or NH when A<sub>2</sub> is alkenyl  
 ending in a double bond;

A<sub>3</sub> is a direct bond, an alkylene or substituted  
 35 alkylene bridge of 1 to 10 carbons, an alkenyl or

- substituted alkenyl bridge of 2 to 10 having one or more double bonds, an alkynyl or substituted alkynyl bridge of 2 to 10 carbons having one or more triple bonds,  
 $-(CH_2)_d-O-(CH_2)_e-$ ,  $-(CH_2)_d-S-(CH_2)_e-$ , or  
 5  $-(CH_2)_d-(C=O)-(CH_2)_e-$ ;  
     d is zero or an integer from 1 to 6;  
     e is zero or an integer from 1 to 6;  
     R<sub>5</sub> is hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, -A<sub>1</sub>-aryl,  
 10 substituted aryl, -A<sub>1</sub>-substituted aryl, heterocyclo, -A<sub>1</sub>-heterocyclo, heteroaryl or -A<sub>1</sub>-heteroaryl;  
     R<sub>9</sub>, R<sub>10</sub>, R<sub>11</sub>, R<sub>12</sub>, R<sub>13</sub>, R<sub>15</sub>, R<sub>16</sub>, and R<sub>19</sub> are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl,  
 15 substituted cycloalkyl, aryl, substituted aryl, heterocyclo, heteroaryl, -A<sub>1</sub>-cycloalkyl, -A<sub>1</sub>-substituted cycloalkyl, -A<sub>1</sub>-aryl, -A<sub>1</sub>-substituted aryl, -A<sub>1</sub>-heterocyclo and -A<sub>1</sub>-heteroaryl, or R<sub>12</sub> and R<sub>13</sub> taken together with the N atom to which they are attached represent a heterocyclo  
 20 ring;



- represents a monocyclic heterocyclo or heteroaryl ring of 4 to 8 atoms containing up to 3 additional heteroatoms (up to 2 additional heteroatoms when the ring is 4 atoms) which are selected from one or  
 25 two oxygen atoms and/or one or two sulfur atoms and/or one, two or three nitrogen atoms;

- R<sub>21</sub> is attached to an available carbon or nitrogen atom and is hydrogen, alkyl, halogen, hydroxy, trifluoromethyl, amino, alkoxy or carboxy;  
 30 R<sub>22</sub> is attached to an available carbon or nitrogen atom and is keto,  $-(C=O)R_{23}$ ,  $-CO_2-R_{23}$ ,  $-NH-(C=O)-R_{23}$ ,  $-N(alkyl)_2$ , -A<sub>1</sub>-hydroxy, -A<sub>1</sub>-N(R<sub>9</sub>)(R<sub>10</sub>), -A<sub>1</sub>-alkoxy, -A<sub>1</sub>-carboxy, -A<sub>2</sub>-cycloalkyl, -A<sub>2</sub>-substituted cycloalkyl, -A<sub>2</sub>-aryl, -A<sub>2</sub>-substituted aryl,


-A<sub>2</sub>-heterocyclo, or -A<sub>2</sub>-heteroaryl;

n is one or two;


m is zero or one;


R<sub>23</sub> is alkyl, -N(R<sub>9</sub>)(R<sub>10</sub>), -A<sub>1</sub>-hydroxy, -A<sub>1</sub>-N(R<sub>9</sub>)(R<sub>10</sub>),

- 5 -A<sub>1</sub>-carboxy, -A<sub>2</sub>-cycloalkyl, -A<sub>2</sub>-substituted cycloalkyl, -A<sub>2</sub>-aryl, -A<sub>2</sub>-substituted aryl, -A<sub>2</sub>-heterocyclo or -A<sub>2</sub>-heteroaryl;




represents a fused bicyclic ring wherein



the monocyclic ring  is defined previously and


- 10  represents a cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclo or heteroaryl having two carbon atoms in common with the monocyclic ring

; and



represents a spiro ring wherein the

- 15 monocyclic ring  is defined previously and  represents a cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclo or heteroaryl ring having a

common carbon with the monocyclic ring .




- 20 2. A compound of Claim 1 including a pharmaceutically acceptable salt thereof wherein:

Y is nitrogen;

Z is nitrogen or C(R<sub>7</sub>);

- 25 E<sub>1</sub> is -O-R<sub>1</sub>, -NH-A<sub>1</sub>-cycloalkyl, -NH-A<sub>1</sub>-heterocyclo, or -NH-A<sub>1</sub>-heteroaryl;

$E_2$  is  $-NH-A_1$ -alkoxy,  $-NH-A_1-N \begin{matrix} R_{15} \\ R_{16} \end{matrix}$ ,  $-NH-A_1$ -substituted phenyl, or  $-NH-A_1-CO_2$ -alkyl;  
 $R_1$  is  $-A_1$ -substituted phenyl;  
 $X_1$  is  $-O-A_1$ -heteroaryl,  $-O-A_1$ -heterocyclo,



10  $R_7$  is hydrogen;

$R_4$  is hydrogen;

$R_3$  is straight or branched chain alkyl of 1 to 4 carbons;

15  $R_5$  is hydrogen, alkyl,  $-CO_2$ -alkyl,  $-A_1$ -phenyl, or  $-A_1$ -heteroaryl wherein alkyl is straight or branched chain of 1 to 4 carbons;

20  $R_2$  is  $-(C=O)R_9$ ,  $-S-R_9$ ,  $-O-R_9$ ,  $-N(R_9)(R_{10})$ ,  $-NR_{11}CO_2R_{19}$ ,  $-(C=O)N(R_{11})CH_2CO_2R_{19}$ ,  $-NR_{11}(C=O)R_{19}$ ,  $-CO_2R_9$ , nitrogen when  $A_2$  is alkynyl ending in a triple bond,  $-(C=O)N(R_{12})(R_{13})$ , phenyl, substituted phenyl, cycloalkyl, heterocyclo, heteroaryl, cycloalkyl- $A_3$ -phenyl, phenyl- $A_3$ -heteroaryl, heterocyclo- $A_3$ -phenyl, or heterocyclo- $A_3$ -heterocyclo;



$R_{25}$  is  $-S-R_9$ ,  $-NR_{11}CO_2R_{19}$ , nitrogen when  $A_2$  is alkynyl ending in a triple bond,  $-(C=O)N(R_{11})CH_2CO_2R_{19}$ ,  $-NR_{11}(C=O)R_{19}$ ,  $-CO_2R_9$ ,  $-(C=O)N(R_{12})(R_{13})$ , phenyl, substituted phenyl, cycloalkyl, heterocyclo, heteroaryl, cycloalkyl- $A_3$ -phenyl, heterocyclo- $A_3$ -phenyl, phenyl- $A_3$ -heteroaryl or heterocyclo- $A_3$ -heterocyclo;

$A_1$  is an alkylene or substituted alkylene bridge of 1 to 6 carbons wherein said substituent is a straight or branched chain alkyl of 1 to 4 carbons;

$A_2$  is a direct bond, an alkylene or substituted alkylene bridge of 1 to 6 carbons wherein said substituent is one or two members selected from alkyl, phenyl, substituted phenyl,  $-CO_2$ -alkyl, carboxy, hydroxy,  $-NH-(C=O)$ -alkyl, and  $-CH_2-(C=O)-NH_2$ , an alkenyl bridge of 2 to 4 carbons having one double bond, or an alkynyl bridge of 2 to 3 carbons having one triple bond wherein alkyl is straight or branched chain of 1 to 4 carbons;

the term "heterocyclo" refers to a substituted or unsubstituted fully saturated or partially saturated 5 to 7 membered monocyclic rings containing one or two heteroatoms selected from oxygen, sulfur and nitrogen and bicyclic rings wherein the monocyclic ring as defined above is fused to a phenyl or substituted phenyl or wherein a bridge of 2 or 3 carbons is present between available carbon and nitrogen atom, said nitrogen and sulfur atoms may optionally be oxidized and said nitrogen atom may optionally be quaternized; said heterocyclo group may be attached at any available nitrogen or carbon atom, and said heterocyclo ring may contain one or two substituents attached to an available carbon or nitrogen atom selected from alkyl, keto and  $-CO_2$ -alkyl, wherein alkyl is straight or branched chain of 1 to 4 carbons;

the term "heteroaryl" refers to a substituted or unsubstituted aromatic 5 or 6 membered monocyclic ring containing one or two oxygen or sulfur atoms and/or from

one to four nitrogen atoms provided that the total number of heteroatoms in the ring is four or less, and bicyclic rings wherein the monocyclic ring as defined above is fused to a phenyl or substituted phenyl, said nitrogen  
 5 and sulfur atoms may optionally be oxidized and said nitrogen atom may optionally be quaternized, said heteroaryl group may be attached at any available nitrogen or carbon atom, and said heteroaryl ring may contain one or two substituents attached to an available  
 10 carbon or nitrogen atom selected from straight or branched chain alkyl of 1 to 4 carbons and halo;

the term "cycloalkyl" refers to a fully saturated cyclic hydrocarbon group of 3 to 7 carbons and such cycloalkyl rings fused to a phenyl ring or such  
 15 cycloalkyl rings 5 to 7 carbons having a carbon-carbon bridge of 3 or 4 carbons;

the term "substituted phenyl" refers to a phenyl ring having one, two, or three substituents selected from alkyl, halo, hydroxy, trifluoromethyl, alkoxy of 1 to 4  
 20 carbons,  $-N(alkyl)_2$ , and  $SO_2NH_2$  wherein alkyl is straight or branched chain of 1 to 4 carbons, and a phenyl ring substituted with a fused five membered ketal;

$R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$ ,  $R_{13}$ ,  $R_{15}$ ,  $R_{16}$  and  $R_{19}$  are independently selected from the group consisting of  
 25 hydrogen, straight or branched chain alkyl of 1 to 4 carbons, phenyl, substituted phenyl and  $-A_1$ -phenyl;

$A_3$  is a direct bond, an alkylene bridge of 1 to 6

carbons, or  $-CH_2-\overset{\overset{O}{\parallel}}{C}-$  ;


30  $-\text{N} \bigcirc$  is a 5 to 7 membered heterocyclo ring which can contain an additional nitrogen atom or can contain an oxygen or sulfur atom;

$R_{21}$  is attached to an available carbon or nitrogen atom and is hydrogen, straight or branched chain alkyl of 1 to 4 carbons, hydroxy or amino;

$R_{22}$  is attached to an available carbon or nitrogen atom and is keto,  $-(C=O)R_{23}$ ,  $-CO_2R_{23}$ ,  $-NH-(C=O)-R_{23}$ ,  
 5  $-N(alkyl)_2$ ,  $-A_1$ -hydroxy,  $-A_1-N(R_9)(R_{10})$ ,  $-A_1$ -alkoxy,  $-A_2$ -phenyl,  $-A_2$ -substituted phenyl, or  $-A_2$ -heteroaryl wherein alkyl is straight or branched chain of 1 to 4 carbons and alkoxy is such an alkyl bonded through an oxygen;

10  $n$  is one or two;  
 $m$  is zero or one;

$R_{23}$  is alkyl,  $-N(R_9)(R_{10})$ , or  $-A_2$ -heteroaryl wherein alkyl is straight or branched chain of 1 to 4 carbons;




represents a fused bicyclic ring wherein

15 the monocyclic ring is defined previously and



represents a substituted phenyl having two carbon

atoms in common with the monocyclic ring ; and




represents a spiro ring wherein the

monocyclic ring is defined previously and   
 20 represents a heterocyclic ring having a common carbon with

the monocyclic ring .


3. A compound of Claim 2 including a pharmaceutically acceptable salt thereof wherein:

25  $Z$  is  $CH$ ;  
 $R_3$  is ethyl;




$E_2$  is -NH-CH<sub>2</sub>-disubstituted phenyl;


- 5 the term "disubstituted phenyl" refers to a phenyl ring having two substituents independently selected from halogen and methoxy or wherein said disubstituted phenyl





- 10  $X_1$  is -O-A<sub>1</sub>-heterocyclo, -O-A<sub>1</sub>-heteroaryl, -NH-A<sub>2</sub>-R<sub>2</sub>,



$X_2$  is  $-O-A_1$ -heterocyclo,  $-O-A_1$ -heteroaryl,  $-NH-A_2-R_{25}$ ,





4. A compound of Claim 3 including a  
5 pharmaceutically acceptable salt thereof wherein:




10

$X_1$  and  $X_2$  are independently selected from the group consisting of










7. The compound of Claim 6 including a pharmaceutically acceptable salt thereof wherein:



5

8. A compound of Claim 1 selected from the group consisting of:

4-[[[4-[[[3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]amino]-1-piperidinecarboxylic acid, ethyl ester;

10



- 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-  
N-[3-(2-oxo-1-pyrrolidiny)propyl]-1H-pyrazolo[3,4-  
b]pyridine-5-carboxamide;
- 5
- 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-  
N-(2-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-  
carboxamide;
- 10
- 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-  
N-(3-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-  
carboxamide;
- 15
- 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-  
N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-  
carboxamide;
- 20
- 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-  
N-[(4-hydroxy-3-methoxyphenyl)methyl]-1H-pyrazolo[3,4-  
b]pyridine-5-carboxamide;
- 25
- 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-  
N-[[4-(1,2,3-thiadiazol-4-yl)phenyl]methyl]-1H-  
pyrazolo[3,4-b]pyridine-5-carboxamide;
- 30
- 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-  
N-(3-1H-imidazol-1-ylpropyl)-1H-pyrazolo[3,4-b]pyridine-  
5-carboxamide;

N-[(3-chloro-4-methoxyphenyl)methyl]-4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;

- 5 (R)- $\alpha$ -[[4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-amino]benzeneacetic acid, ethyl ester;

- 10 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[1-(phenylmethyl)-4-piperidinyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;

- 15 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[3-(4-morpholinyl)propyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;

- 20 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[(4-pyridinyl-1-oxide)methyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;

4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(2-furanylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;

- 25 N-[[4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-L-valine, ethyl ester;

- 30 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(1,2,3,4-tetrahydro-1-naphthylenyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;

- 35 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(tetrahydro-2-furanylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;

- 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-N-[[4-(dimethylamino)phenyl]methyl]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;
- 5 N-[[4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-L-alanine, methyl ester;
- 10 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[[4-fluoro-2-(trifluoromethyl)phenyl]methyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;
- 15 N-[[4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-L-methionine, methyl ester;
- 20 2-[[[4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-amino]propanedioic acid, diethyl ester;
- 25 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(2-oxo-2-phenylethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;
- 30 4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[2-(4-morpholinyl)ethyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;
- 35 N-[[4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-yl]carbonyl]-L-asparagine, 1,1-dimethylethyl ester;
- (S)-4-[[ (3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[2-oxo-3-[2-oxo-2-(1-pyrrolidinyl)-

ethyl]cycloheptyl]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;

5 (S)- $\alpha$ -[[[4-[(3-chloro-4-methoxyphenyl)methyl]-amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-yl]carbonyl]amino]-4-hydroxybenzeneacetic acid;

10 4-[(3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(2-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridin-5-carboxylate;

15 4-[(3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridin-5-carboxylate;

4-[(3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-[3-(4-morpholinyl)ethyl]-1H-pyrazolo[3,4-b]pyridin-5-carboxylate;

20 4-[(3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(4-pyridinylethyl)-1H-pyrazolo[3,4-b]pyridin-5-carboxamide;

25 4-[(3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(2-pyridinylethyl)-1H-pyrazolo[3,4-b]pyridin-5-carboxamide;

30 (R)-4-[(3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(2-tetrahydrofurylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;

35 (S)-4-[(3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(2-tetrahydrofurylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;

4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-N-(hexahydro-2-oxo-1H-azepin-3-yl)-1H-pyrazolo[3,4-b]pyridin-5-carboxamide;

5 4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-N-(cyanomethyl)-1-ethyl-1H-pyrazolo[3,4-b]pyridin-5-carboxamide;

10 (S)- $\alpha$ -[[[4-[[[(3-chloro-4-methoxyphenyl)-methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-yl]carbonyl]amino-4-hydroxybenzeneacetic acid;

15 4-[[[4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-1-ethyl-1H-pyrazolo[3,4-b]pyridine-5-yl]carbonyl]amino]butanoic acid, methyl ester; and;

20 4-[[[(3-chloro-4-methoxyphenyl)methyl]hydroxyl]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridin-5-carboxamide or a pharmaceutically acceptable salt of any of the above compounds.

9. A pharmaceutical composition useful for treating a cGMP associated condition comprising a compound of Claim 1 including a pharmaceutically  
25 acceptable salt thereof and a pharmaceutically acceptable carrier.


10. A pharmaceutical composition useful for treating a cGMP associated condition comprising a  
30 compound of Claim 8 including a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

11. The pharmaceutical composition of Claim 10  
35 wherein the compound is 4-[[[(3-chloro-4-

methoxyphenyl)methyl]amino]-1-ethyl-N-(4-pyridinylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide or a pharmaceutically acceptable salt thereof.

- 5 12. A pharmaceutical composition useful for treating a cGMP associated condition comprising a pharmaceutically acceptable carrier and a compound of the formula

10 (III)



including a pharmaceutically acceptable salt thereof wherein  $E_2$ , Y, Z,  $R_3$  and  $R_4$  are as defined in Claim 1;

$X_3$  is  $-O-R_9$ ,  $-O-A_1-O-R_9$ ,  $-N(R_9)(R_{10})$ ,  $\begin{array}{c} -N-A_2-O-R_9 \\ | \\ R_5 \end{array}$ ,


15  $\begin{array}{c} -N-A_1-N(R_9)(R_{10}) \\ | \\ R_5 \end{array}$ , or a monocyclic ring  $\begin{array}{c} -N \\ | \\ (R_{21})_n \end{array}$ ; and

$R_9$ ,  $R_{10}$ ,  $R_5$ ,  $A_1$ ,  $A_2$ , and  $\begin{array}{c} -N \\ | \\ (R_{21})_n \end{array}$  are as defined in

Claim 1.

- 20 13. The composition of Claim 12 wherein:  
Y is nitrogen;  
Z is CH;  
25  $R_3$  is ethyl;  
 $R_4$  is hydrogen;

25



14. A method of treating a mammalian host to  
 5 relieve a cGMP associated condition comprising  
 administering to said host an effective amount of a  
 composition of Claim 9.

15 15. The method of Claim 14 wherein the cGMP  
 associated condition being treated is erectile  
 dysfunction.

16. A method of treating a mammalian host to  
 relieve a cGMP associated condition comprising  
 15 administering to said host an effective amount of a  
 composition of Claim 10.

17. The method of Claim 16 wherein the cGMP  
 associated condition being treated is erectile  
 20 dysfunction.

18. A method of treating a mammalian host to  
 relieve a cGMP associated condition comprising  
 administering to said host an effective amount of a  
 25 composition of Claim 12.

19. A method of Claim 18 wherein the cGMP  
 associated condition being treated is erectile  
 dysfunction.

30

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/21070

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : A61K 31/415, 31/41; C07D 471/02

US CL : 514/303, 300; 546/119, 113

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/303, 300; 546/119, 113

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS Online, Registry, USPatFull, WEST

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 116:143597, OQUIST, N.L. 'Inotropic responses to selective (RO 20-1724 and SQ 65,442) and nonselective (trequinsin) inhibitors of cyclic AMP-specific class IV phosphodiesterase in newborn, immature, and adult rabbit myocardium,' abstract, <i>Pediatr. Res.</i> , 1992.	1-19
A	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 113:207309, TOMES, C. 'cAMP analogs and selective inhibitors used to study low Km Mucor rouxii cAMP phosphodiesterase,' abstract, <i>Int. J. Biochem.</i> , 1990.	1-19

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier document published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

03 NOVEMBER 1999

Date of mailing of the international search report

02 FEB 2000

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

D. MARGARET SEAMAN

Telephone No. (703) 308-1235



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/21070

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Database CA on STN, Chemical Abstracts, (Columbus OH, USA), No. 112:441, AHN, H.S. 'Effects of selective inhibitors on cyclic nucleotide phosphodiesterases of rabbit aorta,' abstract, Biochem. Pharmacol., 1989.	1-19
A	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 108:48712, DAVIS, A. 'Strategic approaches to drug design,' abstract, J. Comput.-Aided Mol. Des., 1987.	1-19
A	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 107:214090, KURTZ, M.K. 'Cyclic nucleotide phosphodiesterases in cultured normal and RCS rat pigment epithelium: kinetics of cyclic AMP and cyclic GMP hydrolysis,' abstract, Exp. Eye Res., 1987.	1-19